Distribution and important biological features of coastal fish resources in Southeast Asia



LIBRARY
JUL 13 1987

FAO FISHERIES TECHNICAL PAPER

278

FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS



Digitized by the Internet Archive
in 2025

https://archive.org/details/fao-fisheries-and-aquaculture-technical-paper_1986_278

Distribution and important biological features of coastal fish resources in Southeast Asia

by
Somsak Chullasorn
Marine Fisheries Division
Department of Fisheries
Bangkok, Thailand
and
Purwito Martosubroto
Research Institute for Marine Fisheries
Jakarta, Indonesia

FAO
FISHERIES
TECHNICAL
PAPER

278

FOOD
AND
AGRICULTURE
ORGANIZATION
OF THE
UNITED NATIONS
Rome, 1986

The designations employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

M-43
ISBN 92-5-102495-2

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying or otherwise, without the prior permission of the copyright owner. Applications for such permission, with a statement of the purpose and extent of the reproduction, should be addressed to the Director, Publications Division, Food and Agriculture Organization of the United Nations, Via delle Terme di Caracalla, 00100 Rome, Italy.

© FAO 1986

PREPARATION OF THIS PAPER

This document has been prepared as part of FAO's Regular Programme activities, and aims to review all the available information on the biological features of the major marine resources in the region where intensive fishing has been carried out and where the establishment of a firm basis for rational management is therefore required. It also focuses on the preparation of geographic maps of important features of the major resources and fisheries to provide visual material for both resource assessment and management purposes.

ACKNOWLEDGEMENT

The authors' sincere thanks are due first to the fisheries authorities and institutes of the Member Nations in the region and to the many scientists concerned for their assistance in collecting valuable information on coastal fish resources and fisheries, and second to the international organizations in the region, including the Southeast Asian Fisheries Development Center (SEAFDEC) in Bangkok, the International Center for Living Aquatic Resources Management (ICLARM) in Manila, and the FAO/UNDP South China Sea Fisheries Development and Coordinating Programme (SCSP), also in Manila, for their cooperation.

Gratitude should also be extended to the participants of the FAO/SEAFDEC Workshop on Shared Stocks in Southeast Asia, held in Bangkok on 18-22 February 1985 (FAO/SEAFDEC, 1985), who assisted in compiling and revising the contents of this study at the earlier stage of the study.

Finally, the authors are pleased to acknowledge the considerable assistance given by many FAO staff members in undertaking this study, especially Dr S. Chikuni, Fishery Resources and Environment Division, for his coordination throughout the study and for the technical editing of the draft manuscript, and to Miss Gloria A. Soave, Research Information Unit, for revising the bibliography at the final stage.

Distribution:

Authors
FAO Fisheries Department
FAO Regional Fisheries Officers
Resource Information/Fish Biology/
Species Identity/Fishery Management

For bibliographic purposes this document should be cited as follows:

Chullasorn, S. and P. Martosubroto, 1986. Distribution and important biological features of coastal fish resources in Southeast Asia. FAO Fish.Tech.Pap., (278): 84 p.

ABSTRACT

The distribution and important biological features of the major coastal fish resources in the region, covering finfish, both pelagic and demersal, shrimps and cephalopods, are reviewed. First, the names of the major species or species group reported from the countries bordering on the region are shown by country in a table. Second, the information and parameters on important biological features of major species are tabulated by species group in subsequent tables with specific studies. Third, the geographic distribution of major species groups is shown on schematic illustrations. The sources of the information for the mapping is given in both the text and the references at the end of this paper. The results obtained are discussed first in a general manner, and second in connection with the requirements for future research.

CONTENTS

		Page
1.	INTRODUCTION	1
2.	SOURCES	1
3.	AREA EMPLOYED AND ENVIRONMENTAL CONDITIONS	2
4.	OCCURRENCE OF MAJOR SPECIES GROUPS AND SPECIES	3
5.	IMPORTANT BIOLOGICAL FEATURES AND PARAMETERS OF MAJOR SPECIES	3
6.	GEOGRAPHIC DISTRIBUTION OF HABITAT, SPAWNING AND FISHING GROUNDS OF MAJOR SPECIES GROUPS	3
	6.1 Coastal Pelagic Fish	3
	6.1.1 Mackerels	4
	6.1.2 Round scads	5
	6.1.3 Sardines	5
	6.1.4 Anchovies	6
	6.1.5 Carangids	6
	6.1.6 King mackerels	7
	6.1.7 Coastal tunas	7
	6.2 Demersal Fish	7
	6.2.1 Pony fish, Leiognathidae	8
	6.2.2 Red snappers, Lutjanidae	9
	6.2.3 Goat fish, Mullidae	9
	6.2.4 Threadfin breams, Nemipteridae	9
	6.2.5 Bigeye snappers, Priacanthidae	10
	6.2.6 Groupers, Serranidae	10
	6.2.7 Croakers, Sciaenidae	10
	6.2.8 Lizardfish, Synodontidae	10
	6.3 Shrimps	11
	6.4 Cephalopods	12
7.	DISCUSSION	12
8.	REFERENCES	14
LIST OF TABLES		27
LIST OF FIGURES		63

1. INTRODUCTION

The waters in Southeast Asia constitute one of the most productive areas for commercial fisheries in the world oceans. The total annual catch from the region has increased steadily throughout the years and has reached about 6 million tons recently, which accounts for about 10 percent of the world total marine production.

Although all nations in the region have traditionally harvested marine resources in their coastal waters, the history and the current status of the fisheries vary greatly from country to country. For instance, trawl fishing was first initiated in Thailand in the early 1960s, followed by Malaysia and Indonesia, while in the Philippines fishing for pelagic fish developed faster. In Kampuchea and Viet Nam, on the other hand, marine fishing has not developed well in recent years, due mostly to political disturbances. In Burma and Brunei, marine fishing has not been particularly intensive until recently. The same is true for East Malaysia, with the exception of shrimp fishing in Sabah. The historical development process and the current status of the fisheries in the region are reviewed in detail in another FAO publication (Shindo, in preparation).

Because of the development of these fisheries, many resources of the coastal waters of the region have become heavily exploited. Some stocks, especially demersal fish resources in nearshore waters, are believed to be over-exploited, and many coastal pelagic fish resources in shallower waters nearly fully exploitated in recent years. Under these circumstances, many countries have established several regulatory measures aiming at both the conservation of their resources and at the resolution or alleviation of the conflicts between fisheries and fishermen. On the other hand, it is believed that several other stocks in the region may not be fully exploited, and increased catches can therefore be expected from those stocks, especially in offshore areas.

In connection with the extension of national jurisdiction to 200 miles, the proper assessment and rational utilization of shared stocks, have recently become significantly important for both research and management purposes in the region. There is no doubt that international cooperation is critically important in the future to ensure better assessment and management of these resources.

Despite the serious state of many coastal resources in the region as mentioned above, explicit assessment has scarcely been made due mostly to the paucity of the biological information on the resources and the lack of adequate catch and effort data. Thus, it is considered to be the right time to review the biological information available so as to identify the gaps between present knowledge and the requirements for the future.

This paper presents a compilation of all available information on the coastal fish resources in the region, covering both important biological features and fisheries. The definition of the sea area employed in this study is given in Figure 1, and the bathymetric conditions of the area in Figure 2. The biological data are presented in both tabulated forms (Tables 1-6) and maps illustrating the geographic distribution of the resources and the migratory/displacement patterns of major species (Figures 3-21). Oceanic resources, such as skipjack and larger tunas, are excluded from this study, although the fishery has been quite intensive in the eastern part of the region in recent years. They are widely distributed in the Pacific Ocean beyond the border of this particular region, and are highly migratory in nature. The biological studies on these fish have been carried out by several tuna-fishing countries and international organizations, and the review of these studies is therefore beyond the scope of this particular study.

It is expected that the research institutes in the countries concerned will identify, from the information provided in this paper, the priority areas for their future research. It is also hoped that the paper will be useful for fisheries administrators in planning the proper management and development of the fisheries in their countries.

2. SOURCES

Data and information have been collected from both published reports and unpublished documents in various institutions and organizations in the region. Some have also been collected orally through interviews during the visit of the authors to national institutes in Hong Kong, Indonesia, Malaysia, the Philippines and Thailand. Visits were also made to several international organizations such as the Southeast Asian Fisheries Development Center (SEAFDEC) in Bangkok, the International Center for Living Aquatic Resources Managment (ICLARM), and FAO/UNDP South China Sea Fisheries Development and Coordinating Programme (SCSP), both in Manila.

It should be noted here that substantial additions and revisions of the information collected by the authors were made by the participants of the FAO/SEAFDEC Workshop on Shared Stocks in Southeast Asia held in Bangkok during 18-22 February 1985 (FAO/SEAFDEC, 1985), where the interim draft paper of this study was submitted for discussion.

The list of all the publications and documents referred to in this review is given in the References at the end of this paper. The specific literature to refer to in each of the species group or species is given in each of the following sections, sub-sections and tables in the paper. However, a few papers that have been used on a global basis throughout the study are given here. These are Morgan and Valencia (1983) and Wyrtki (1961) for resources/fisheries/environment in Southeast Asia, Marr (1976) and Tiews (1976) for resources/management in Southeast Asia, Shindo and Chullasorn (1980) for resources/fisheries in Southeast Asia, SEAFDEC (1984) for fishery statistics in Southeast Asia, FAO/SEAFDEC (1985) for shared stocks in Southeast Asia, Beales et al. (1982) for resources/fisheries in Brunei, Blanc (1960) and FAO (1980) for resources/fisheries in Kampuchea, Burma (1982) and Khin (1948) for resources/fisheries in Burma.

3. AREA EMPLOYED AND ENVIRONMENTAL CONDITIONS (Figures 1 and 2)

The marine waters covered here fall entirely in the tropical zone of Southeast Asia. They are composed of several major seas and straits (Figure 1), namely the Andaman Sea in the west, the Strait of Malacca, the South China Sea, the Java Sea, the Arafura Sea, the other Indonesian waters including the Makassar Strait, the Molucca, Flores, Savu, Timor and Banda Seas, the Indian Ocean along the Indonesian archipelago in the south, the Celebes Sea, the Sulu Sea, the other Philippino waters such as the Sibuyan, Visayan and Mindanao Seas, the Pacific Ocean along the Philippines to Indonesia coasts in the east and the Formosa Strait in the north, which connects the region to the East China Sea further north.

Twelve countries bordering on the region are all traditionally fishing nations though the magnitude of fishing effort varies from one to the other (Figure 1). They are Brunei, Burma, China including both the mainland and Taiwan Province, Hong Kong, Indonesia, Kampuchea, Macau, Malaysia, Philippines, Singapore, Thailand and Viet Nam.

Bathymetric conditions divide the region roughly into two parts, namely shallow and flat areas in the west where wide continental shelves have been well developed and deep sea areas in the east with narrow continental shelves and steep slopes (Figure 2). The former include the Strait of Malacca, the southern part of the South China Sea including the Gulf of Thailand, the Gulf of Tong King, the northwestern part of the South China Sea, the Java and Arafura Seas, while the rest of the region are covered by the latter. These characteristics principally determine the occurrence of exploitable resources. For instance, there are abundant demersal fish and shrimp/prawn resources in the former areas and oceanic fish resources such as skipjack and larger tunas in the latter.

The climatic and oceanographic conditions of the region are generally governed by the tropical monsoon regime (Wyrtki, 1961; Marr, 1976). The waters are also strongly influenced by the inflow from the neighbouring seas and oceans in accordance with the change in monsoons. For instance, during the northeast monsoon the water from the Pacific Ocean and the East China Sea move into the area through the northern part of the South China Sea, while during the southwest monsoon from both the Pacific and Indian Oceans through the southern part of the region, although the magnitude is smaller than the former.

The surface sea current in the region changes its direction completely in accordance with the two monsoon seasons, with the exception of the Malacca Strait where waters flow always in a northerly direction, although the intensity varies according to the season. However, seasonal changes in water temperature are very slight. The temperature usually ranges between 24° and 29°C throughout the year, with the exception of the northern part of the South China Sea where it decreases to below 24°C in winter due mostly to the inflow of cold waters from the East China Sea.

These are the distinctions in the environmental characteristics of the region that determine the species and ecological features of the resources. The region-specific features of the resources, which have been derived from the above-mentioned environment in the region, are the multispecies nature of fish community; the fast growing and early maturing of the individual fish with relatively short life span; and the multiple or almost all year-round prolonged spawning.

The region is also characterized by the abundant distribution of coral reefs, especially in the western-most and eastern parts, while mangroves are numerous where there is substantial discharge of fresh water. These environments also play an important role in enhancing the productivity of the coastal waters, providing favourable conditions for spawning and nursing of many fish and shrimp resources.

4. OCCURRENCE OF MAJOR SPECIES GROUPS AND SPECIES (Table 1)

The list of the major species groups and species reported to have been found in each of the national jurisdictional waters is given in Table 1. For pelagic fish, 46 groups have been identified and 29 species of commercial value have been confirmed to be commonly distributed; for demersal fish, 39 groups and 36 species; for shrimps, 8 groups and 13 species; and for squids/ cuttlefish, 8 groups and 6 species. The information in the table was collected from the many references mentioned in the other sections and tables in this paper.

It should be noted, however, that the checklist shown in Table 1 is still incomplete for many species in some local waters as there has been insufficient research and study on the identification of fish species. Therefore, the blank columns in Table 1 do not mean the lack of species in that specific water but should be read as it being uncertain whether the species really occurs or not. The situation is particularly weak for squids and cuttlefish, which may be the result of the little research made so far due to the unpopularity of these resources in many countries, with the exception of Hong Kong and Thailand.

Despite the incompleteness of the above checklist, the table as a whole gives a clear picture of the multispecies nature of the major resources in the region with a number of species groups and species in each of the fish type categories.

5. IMPORTANT BIOLOGICAL FEATURES AND PARAMETERS OF MAJOR SPECIES (Tables 2 to 6)

The data and information on the important biological features of the major species are given in the subsequent tables, together with the substantive studies to be referred to for each species. They comprise 24 species of coastal small pelagic fish (Table 2), a king mackerel and three species of coastal tunas (Table 3), 42 species of demersal fish (Table 4), 18 species of shrimps (Table 5), and 11 species of cephalopods (Table 6). Some parameters vary according to the localities where sample specimens were taken, which may derive partly from the difference in ecological features caused by the various environmental conditions (e.g., spawning and growth) and partly from the magnitude of fishing intensity (e.g., mortality). For this reason, the results of the studies are given in these tables by country, locality and occasionally even by the scientist involved.

The data appear to be plentiful when taking a glance of the tables. However, it should be noted that their quantity and quality vary greatly according to the area surveyed. Such an imbalance in knowledge in the region is commonly observed in many species but is particularly serious for some specific species groups, e.g., coastal tunas and cephalopods.

Both the pelagic and demersal fish resources in Southeast Asia have become very important for commercial fisheries in many countries during the past two decades. The fisheries research institutes in those countries initiated intensive studies on their resources and have carried out quite a lot during the past twenty years. These activities have covered a number of important resources including two species of mackerels, three of round scads, four of anchovies, three of sardines, one of king mackerels, two of pony fish, two of threadfin breams, one of bigeye snappers and two of lizard fish. Apart from these species groups, information on the biology and ecology of shrimps and cephalopods is still limited. More intensified research and study are awaited in the immediate future for these resources.

6. GEOGRAPHIC DISTRIBUTION OF HABITAT, SPAWNING AND FISHING GROUNDS OF MAJOR SPECIES GROUPS (Figures 3 to 21)

The information on the distribution of resources, fishing grounds, concentration of abundance, spawning grounds and estimated migration pattern are shown in Figures 3-21 for the selected species groups of commercial value.

Several criteria were employed in selecting the species group to be illustrated in these figures, based mostly on the fact that they are important species for commercial fisheries, i.e., the larger quantity of the catch taken from the sea, the higher market price and the higher utility for foreign trade. The others included in the mapping are not being intensively or widely exploited at present; however, they are believed to have a large potential harvest for future exploitation.

The following sub-sections give annotations and interpretations for each of the figures provided in this study.

6.1 Coastal Pelagic Fish

The exploitation of coastal pelagic fish resources started a long time ago in Southeast Asia, before that of demersal fish resources. However, the number of species harvested has been considerably smaller than that of demersal fish. The most important species of commercial value are

commonly found throughout the region, and include such genera as Rastrelliger, Decapterus, Sardinella, Stolephorus, Scomberomorus, several genera of carangids other than Decapterus and coastal tunas mainly composed of Euthynnus, Auxis and Thunnus tonggol. Mapping of the geographic distribution has been made for the seven major species groups: mackerels (Figure 3), round scads (Figure 4), sardines (Figure 5), anchovies (Figure 6), carangids (Figure 7), king mackerels (Figure 8) and coastal tunas (Figure 9).

The literature referred to in each of these species groups is given in each of the sub-sections. However, the following literatures have been used on a broader basis covering multiple species groups.

(1) Resources assessment:

SCSP (1978a) for South China Sea and SEAFDEC (1982) for Southeast Asia with emphasis on shared stocks.

(2) Resources and fisheries:

Menasveta, Shindo and Chullasorn (1973) for South China Sea, Boonragsa (1984) for the Andaman Sea, SCSP (1976, 1976a) for the Malacca Strait, Department of Fisheries, Thailand (1984), Menasveta (1980) and Supongpan et al. (1982) for the Gulf of Thailand, Fisheries Research Institute, Malaysia (1984) for Malaysia, Departemen Pertanian, Indonesia (1983) for Indonesia, SCSP (1979) for the Java Sea, Sujastani (1983) for Sumatra, Uktolseja et al. (1979) for Sulawesi, SCSP (1976b) for the Visayan and Sibuyan Seas, SCSP (1977) for the Sulu Sea, the Bohol Sea and the Moro Gulf, SCSP (1978b) for the Pacific coast of the Philippines, SCSP (1979a) for the north and west coasts of Luzon, and Chung and Chiu (1980) for Hong Kong.

(3) Resources research:

Aglen et al. (1981) for Malaysia and FAO (1975) for Viet Nam.

(4) Resources and biology (general):

Dwiponggo and Pauly (in press) for Indonesia and Ingles and Pauly (1984) for the Philippines.

(5) Larval survey:

Vattanachai (1974) for South China Sea, Chayakul et al. (1983) for the Gulf of Thailand and Ordonez et al. (1973) for the Philippines.

6.1.1 Mackerels (Figure 3)

There are three species of mackerels commonly found in the region (Table 1): Indo-Pacific mackerel or short mackerel (Rastrelliger brachysoma), Indian mackerel (R. kanagurta) and island mackerel (R. faughni). The first species is abundant in the coastal waters, while the second and the third are distributed more in offshore waters throughout the region. They are caught mainly by purse seine and gillnet, and occasionally by trawl. The fishing ground extends widely over the coastal waters in the region (Figure 3). It is noted that these three species are generally combined in fishery statistics in many countries as fishermen usually do not separate their catches for marketing.

Extensive studies on the biology and ecology of mackerels have been carried out for R. brachysoma and R. kanagurta in many waters, including the Gulf of Thailand, the Mergui Archipelago, the Andaman Sea, the Malaysian waters, the Java Sea and the Philippine waters. The biological features and parameters of mackerels are shown in Table 2, and their seasonal migratory patterns estimated from tagging experiments are given in Figure 21. The result of tagging also indicated that there may be a fairly long-range movement of R. brachysoma from the southern part of the west coast of Thailand to the west coast of the Malaysian Peninsula. This supports the hypothesis that the mackerel along the Andaman Sea coast is composed of a single stock being shared by the two or more countries bordering the sea.

In addition to the references already specified in Sections 2 and 6.1, the following literature refers specifically to mackerels in the region: Collette et al. (1983) for the entire region, SCSP (1978) for the South China Sea, Boonragsa et al., (1984), Bussarawitch (1984, 1984a, 1984b), and Druzhinin (1968) for the Andaman Sea, Chong and Chua (1974), FAO/UNDP Bay of Bengal Project (1985) and Pathansali (1961, 1967) for the Malacca Strait, Boonprakob (1965, 1967, 1972), Chullasorn (1979), Dhebtaranon and Chotiyaputta (1974), Somjaiwong et al. (1970), Suchondhamarn et al. (1970), Suvapepun and Suwanrumpha (1970), Tabtimtai (1968), Tantiswetratana (1979) and

Vanichkul and Hongskul (1965) for the Gulf of Thailand, Chee (1980) for Malaysia, Gafa (1982) for the Makassar Strait, Sujastani (1974) for the Java Sea and Tan (1970) for the Manila Bay.

6.1.2 Round scads[1/] (Figure 4)

Round scads in the region are represented by eight species of the genus Decapterus (Table 1). Among them, shortfin scad (D. macrosoma) and white tip scad (D. maruadsi) are the most common in many waters in the region. In Burmese waters Indian scad (D. russelli) is considered the most abundant species. However, there has been much confusion between D. russelli and D. maruadsi in many of the countries. It is impossible to examine the biological features for these two species further in detail unless the problem of species identification is sorted throughout the entire region. Under these circumstances, the catch statistics for round scads have generally been compiled on a species-combined basis. Similarly, for the other five minor species, the establishment of species identification and the confirmation of their distribution in this particular region are urgently awaited from a biological point of view. They include stout redtail scad (D. kurroides), mackerel scad (D. macarellus), amberstripe scad (D. muroadsi), temperate scad (D. scombrinus) and slender redtail scad (D. tabl).

Round scads are widely distributed in the region as is shown in Figure 4. They are caught by various types of purse seines, bagnets and gillnets in the offshore waters. There is a possibility of expanding the fishing on round scads to many parts of the region where it has not been intense and where the resources are believed to be abundant.

Biological information on round scads is geographically limited, i.e., the Palawan waters, the Gulf of Thailand and the Java Sea (Table 2). A recent workshop on shared stocks organized jointly by FAO and SEAFDEC recommended, in the light of the importance of these resources, that further research and study on this species group be carried out in the immediate future to meet assessment and management purposes.

The following publications refer specifically to round scads in the region: SCSP (1978) for the South China Sea, Wongchitsue (1984) for the Andaman Sea, FAO/UNDP Bay of Bengal Project (1985) for the Malacca Strait, Chantarasri (1980), Cheunpan (1981), Chullasorn and Yusukswad (1977) and Piyathirathitivorakul (1983a) for the Gulf of Thailand, Atmadja (1982) for the Arafura Sea, Sadhotomo (1983) for the Java Sea, Magnusson (1970), Ronquillo (1974) and Tiews, Ronquillo and Caces-Borja (1970) for the Philippines. See Sections 2 and 6.1 for global and broader references.

6.1.3 Sardines[2/] (Figure 5)

Although sardines in the region are composed of various genera of clupeoids, the fishery has depended mostly on the two genera, Sardinella and Amblygaster. There are 11 species reported from the region for these genera, but the major ones are confined to about five (Table 1).

Among these, fringescale sardinella (S. fimbriata), goldstripe sardinella (S. gibbosa) and spotted sardinella (A. sirm = S. sirm)[3/] are commonly found in almost all the countries in the region. It is interesting to see that the Indian oil sardine (S. longiceps) predominates in the sardine catch in the Bali Strait. In general, sardines are widely distributed in the coastal waters of the entire region (Figure 5). The major fishing gear employed are purse seines, gillnets and bagnets.

Biological studies on sardines have been carried out only for selected waters in the region, namely the waters around the Palawan, the Manila Bay, the Java Sea, the Bali and Malacca Straits and the Gulf of Thailand (Table 2).

The following literature refers specifically to sardines in the region: Whitehead (1985) for the entire region, Chullasorn (1979a) for the Gulf of Thailand, Dwiponggo (1974), Dwiponggo and Uktolseya (1974), Ritterbush (1975) and Sujastani and Nurhakim (1982) for the Bari Strait and Ronquillo (1960) for the Philippines. See Sections 2 and 6.1 for global and broader references.

1/ The English names employed in this sub-section are those used in FAO species catalogue. Carangids of the world, by W. Smith-Vaniz (in preparation)

2/ The English and scientific names employed in this sub-section refer to those defined in the FAO species catalogue on clupeoid fishes of the world recently published (Whitehead, 1985)

3/ See Whitehead (1985)

6.1.4 Anchovies (Figure 6)

Anchovies are commonly distributed in the coastal waters of the region. The fish prefer a neritic environment and concentrate in nearshore waters, including estuarine areas. They are generally composed of several genera: Coilia, Setipinna, Thryssa, Thrissina and Stolephorus. Among 13 species of Stolephorus found in the region, shorthead anchovy (S. heterolobus), and Indian anchovy (S. indicus) are considered the most important. Anchovies are highly abundant along northern Sumatra, Peninsular Malaysia, the east coast of Thailand, Kampuchea and Viet Nam. The fishing grounds are generally confined to the inshore waters along these coasts (Figure 6). The major fishing gear employed are liftnets, bagnets and small-scale purse seines with small mesh size.

Intensive studies have been made mostly on the genus Stolephorus in the Manila Bay, the Java Sea, the Strait of Singapore, the Malacca Strait and the Gulf of Thailand. The species studied so far include spotty-face anchovy (S. bataviensis[1/]), Buccaneer anchovy (S. buccaneeri[2/]), Commerson's anchovy (S. commersonii), shorthead anchovy (S. heterolobus[3/]), Indian anchovy (S. indicus), Hardenberg's anchovy (S. insularis and S. zollingeri[4/])(Table 2). However, a firm basis has not yet been established for the identification of several species as is shown in the footnote of this sub-section. Further study on this aspect, as well as other biological features of the fish, is urgently required.

The following literature refers specifically to anchovies in the region: Whitehead (in preparation) for the entire region, Tham (1974) for the South China Sea, Dhebtaranon (1973), Isara (1972), Sitthichokpun (1970), Supongpan et al. (1984) and Taweesith (1979) for the Gulf of Thailand, Tham (1953, 1967, 1967a) for the Singapore Strait and Tiews, Ronquillo and Santos (1970) for the Manila Bay. See Sections 2 and 6.1 for global and broader references.

6.1.5 Carangids[5/] (Figure 7)

The carangids (excluding Decapterus spp., which have already been dealt with separately) are generally composed of a large number of genera and species in this region, 19 and 49 respectively. Among these, Atule, Carangoides, Megalaspis, Scomberoides, Selar and Selaroides are the most important groups of commercial value. Due to the involvement of so many species, the mixed catches and the difficulties in field identification of species, fishery statistics for this group have generally been compiled at the species combined level in many countries. Torpedo scad[6/] (Megalaspis cordyla) is the only species for which statistics have been separately compiled in all the countries concerned. In the geographic distribution of this group reviewed here, however, all carangid species have been combined and are shown in Figure 7.

Under these circumstances, the biological research carried out so far has been limited in terms of both the species examined and the areas surveyed, and applies to yellowtail scad (Atule mate), cleftbelly trevally[7/] (Atropus atropus), malabar trevally[8/] (Carangoides malabaricus), rainbow runner (Elagatis bipinnulata), torpedo scad[6/] (Megalaspis cordyla), bigeye scad (Selar crumenophthalmus) and yellowstripe scad[9/] (Selaroides leptolepis) in the Gulf of Thailand, Indonesia and Philippine waters (Table 2).

1/ Recently classified as, Stolephorus waitei[10/]

2/ Recently classified as, Encrasicholina punctifer[10/]

3/ Recently classified as, Encrasicholina heteroloba[10/]

4/ Recently classified as, Encrasicholina punctifer[10/], a synonym of the above. English name is therefore not given in this paper.

5/ The English names employed in this sub-section are tentative as FAO standard names (in preparation)

6/ Hardtail scad at present

7/ Kuweh trevally at present

8/ Malabar cavalla at present

9/ Yellowstripe trevally at present

10/ Reference, FAO species catalogue on clupeoid fishes of the world FAO Fish.Synop., (125)Vol. 7, Pt. 2 (Whitehead, in preparation)

The following literature refers specifically to carangids in the region: Termvidchakorn (1983) and Tu (1971) for the South China Sea, Bhatia et al. (1979) and Druzhinin and Hliang (1972) for the Andaman Sea, Latiff et al. (1974) and Wahyuono et al. (1983) for the Malacca Strait, Naiyanetr (1963), Nupetch (1984), Nupetch et al. (1983) and Piyathirathitivorakul (1983) for the Gulf of Thailand, Sudradjat and Nugroho (1983) for the Sunda Shelf and Arce (1981) for the Visayan Sea. See Sections 2 and 6.1 for global and broader references.

6.1.6 King mackerels (Figure 8)

In this group there are three species commonly found in the region: narrow-banded Spanish mackerel (Scomberomorus commerson), Indo-Pacific king mackerel (S. guttatus) and streaked seerfish (S. lineolatus). They are generally widely distributed in both inshore and offshore waters (Figure 8). The juvenile and small fish are often caught by trawl and bamboo stake trap in coastal waters while large fish are caught mainly by gillnet, troll and longline in offshore waters. Although the fish fetch a high market price due to the high demand in many of the countries, the quantity of the catch has remained rather small compared to the other fish groups in pelagic fish resources.

The biological research carried out so far has been limited to certain areas such as the Gulf of Thailand and the Visayan Sea (Table 2). It is believed that the fish migrate widely beyond the borders of national jurisdiction in both nearshore and offshore areas.

The following literature refers specifically to king mackerels in the region: Collette and Nauen (1983) for the entire region, Menasveta, Shindo and Chullasorn (1973) for the South China Sea, Boonragsa (1984) for the Andaman Sea, Chayakul et al. (1983), Chullasorn, Chatiyaputta and Chayakul (1973), Klinmuang (1979), Supongpan (1979) and Supongpan and Chayakul (1979) for the Gulf of Thailand, Uktolseja et al. (1979) for Indonesia, Chung and Chiu (1980) for Hong Kong and Ingles and Pauly (1984) for the Philippines. See Section 2 for global references.

6.1.7 Coastal tunas (Figure 9)

The coastal tunas in the region include mostly kawakawa (Euthynnus affinis), frigate tuna (Auxis thazard) and longtail tuna (Thunnus tonggol). They are abundant and are widely distributed almost over the whole region (Figure 9). Until recently, these fish were caught only incidentally in the waters around the Philippines, Indonesia, Malaysia and Thailand. Being stimulated by the strong demand for the canning industries, which have developed rapidly in recent years, fishing on coastal tunas has become one of the important commercial fisheries in these waters. Although the resources are believed to be abundant over the entire region, fishing has not been well developed in the other areas. Further expansion is therefore quite feasible in many coastal and offshore waters of the region.

Knowledge on the biology of these species is limited mostly to the Gulf of Thailand and partly to the Philippine waters (Table 3).

The following literature refers specifically to coastal tunas in the region: Collette and Nauen (1983) and Uchida (1981) for the entire region, Kikawa and Ferraro (1967) for Indian Ocean, Kume (1973) for the South China Sea, Bhatia and Boonragsa (1980), Boonragsa (1984) and Yesaki (1982) for the Andaman Sea, Chayakul et al. (1983), Chiampreecha (1978), Cheunpan (1984), Klinmuang (1978, 1981) for the Gulf of Thailand, Dwiponggo and Pauly (in press) and Uktolseja et al. (1979) for Indonesia, Ingles and Pauly (1984) and Ronquillo (1963) for the Philippines. See Section 2 for global references.

6.2 Demersal fish

Demersal fish in Southeast Asia comprise numerous species and many of them have been commercially harvested. For instance, along the Thai and the Malaysian coasts of the Gulf of Thailand, approximately 30 families of demersal fish exist, of which more than 300 species have been commercially utilized (Wongratana, 1968).

It is a distinctive feature of the demersal fisheries in the region that a substantial amount of small-size fish of no commercial value are usually caught together with larger fish of commercial value (Meemeskul, 1982; Shinoda, 1978). It is usually dealt with as "trash fish" by the fishermen, and accounts sometimes for more than 50 percent of the total catch, especially in the Gulf of Thailand. Trash fish comprise three different species groups: non-edible species unacceptable by the fish market, edible species of low commercial value and juveniles of commercially important larger species. The third category is of importance for assessment and management purposes as it can account for more than 30% of the total trash fish catch (Shinoda, Lim and Tan 1978). However, this involves many highly complicated problems in conjunction with the multispecies nature of the resources (Shinoda et al., 1979), and no immediate solution can be

expected, as discussed in the later part of this paper. Trash fish are generally discarded at sea immediately after sorting out on board. In a few countries, they are processed as fish meal for animal feed and as fish sauce for human consumption.

Demersal fish are generally abundant on the shallower part of continental shelves with a depth range of 10-60 m. Trash fish generally occur on further shallower shelves between 10 and 30 m deep.

The mapping of the geographic distribution of demersal fish has been made for the eight major species groups commonly found in the region: pony fish (Leiognathidae, Figure 10), red snappers (Lutjanidae, Figure 11), goat fish (Mullidae, Figure 12), threadfin breams (Nemipteridae, Figure 13), bigeye snappers (Priacanthidae, Figure 14), groupers (Serranidae, Figure 15), croakers (Sciaenidae, Figure 16) and lizardfish (Synodontidae, Figure 17).

In addition to the references mentioned in the following sub-sections on each of the species groups, a number of publications refers to multiple species groups on a broader basis. See also Section 2 for global references.

(1) Resource assessment:

SCSP (1978a) for the South China Sea and SCSP (1976, 1976a) for the Malacca Strait.

(2) Multispecies resources assessment:

Pauly and Murphy (1982) for the tropics in general, Pauly (1979) for Southeast Asia, Shinoda et al. (1979) for the South China Sea with special emphasis on the selectivity of mesh-size, and Ritragsa (1976) for the Gulf of Thailand.

(3) Resources and fisheries (general):

Aoyama (1973), Shindo (1973) for the South China Sea, Druzhinin and Hliang (1972) for Burma, Chantawong et al. (1984) for the Andaman Sea, Wahyuono et al. (1983) for the Malacca Strait, Fisheries Research Institute, Malaysia (1984) for Malaysia, Chua and Mathias (1975) for West Sabah, Malaysia, Isarankura (1968) and Menasveta (1980) for the Gulf of Thailand, SCSP (1977a, 1977b), for the Sunda Shelf, Departmen Pertanian, Indonesia (1983) for Indonesia, Sujastani (1983) for Sumatra, SCSP (1979) for the Java Sea, Lai-shing (1968) for Hong Kong, SCSP (1976b) for the Visayan and Sibuyan Seas, SCSP (1977) for the Sulu Sea, the Bohol Sea and the Moro Gulf, SCSP (1978b) for the Pacific coast of the Philippines and SCSP (1979a) for West Luzon.

(4) Resources Research:

Kyokuyo Co.Ltd. (1972) for the South China Sea, Latiff et al. (1974), Pathansali et al. (1974) and Pong (1981) for the Malacca Strait, Aglen et al. (1981) for Malaysia, Pong et al. (1976) for West Sarawak, Wongratana (1968) for the Gulf of Thailand, Saeger et al. (1976) and Widodo (1976) for Indonesia.

(5) Resources and biology (general):

Nguyen (1972) and Tu (1971) for the South China Sea, Meemeskul (1978), Pramokechutima (1978) and Ritragsa, Dhamniyon and Sitthichaikasem (1968) for the Gulf of Thailand, Dwiponggo and Pauly (in press) for Indonesia, Martosubroto (1982) for the Java Sea, and Ingles and Pauly (1984) for the Philippines.

(6) Resources/fisheries/biology of trash fish:

Shinoda, Lim and Tan (1978), Shinoda et al. (1979) for the South China Sea and Meemeskul (1982) for the Gulf of Thailand.

6.2.1 Pony fish, Leiognathidae (Figure 10)

There are two genera in this group occurring in the region: Leiognathus and Gazza. The former genus predominates over the latter in terms of the number of species involved as it consists of about 20 species and their abundance. Among them L. bindus, L. blochii, L. brevirostris, L. equulus and L. splendens are very common in the region. They occur in large quantity in shallow waters along the coast throughout the region and are caught mainly by trawl (Figure 10). In the Java Sea the catch of L. splendens accounts for the largest component of the total catch by liftnet.

Biological studies have been carried out for several species of Leiognathus in various waters such as the Manila Bay, the Samar Sea, the Java Sea, the Malacca Strait and the Gulf of Thailand. Results of these studies are shown in Table 4.

The following literature refers specifically to pony fish in the region: Amaragsa (1968) and Vadhanakul (1976) for the Gulf of Thailand and Tiews et al. (1972) for Manila Bay and San Miguel Bay. See Sections 2 and 6.2 for global and broader references.

6.2.2 Red snappers, Lutjanidae (Figure 11)

There are four genera occurring in the region: Lutjanus, Pristipomoides, Caesio[1/] and Glabrilutjanus[2/]. However, Lutjanus is the most important as it is composed of 24 species. Among those species L. argentimaculatus, L. johni, L. lineolatus[3/], L. malabaricus, L. sebae, L. sanguineus and L. vitta are commonly found in commercial catches. They are distributed widely over the areas but a larger concentration is found in offshore waters in the South China Sea deeper than 60 m (Figure 11). There may be similar concentrations in deeper waters in other areas where further expansion of fishing would be feasible. This may be most likely on the deeper and untrawlable ground, where demersal resources have never been heavily fished, if suitable fishing methods are introduced.

Biological features have been studied for four species collected from the Strait of Malacca, the South China Sea (Hong Kong, Malaysia, Singapore), the Gulf of Thailand, the Andaman Sea and the northern Java Sea (Table 4).

The following literature refers specifically to red snappers in the region: Allen (1985) for the entire region, Kangvankij (1970) for the South China Sea, Intong (1980) for the Gulf of Thailand, and Gafa and Sadhotomo (1983) for South Kalimantan. See Sections 2 and 6.2 for global and broader references.

6.2.3 Goat fish, Mullidae (Figure 12)

Goat fish in the region are composed of three genera and 14 species. The most common genus is Upeneus, in which three species are commonly found in commercial catches: U. bensasi, U. moluccensis and U. sulphureus. They are distributed widely in both the coastal and offshore waters. However, they tend to be more abundant in deeper waters. The geographic distribution is given in Figure 12.

Knowledge on the biological features of the fish is limited to a few specific areas: the South China Sea (Hong Kong), the Java Sea and the Philippine waters (Table 4).

The following literature refers specifically to goat fish in the region: Badrudin (1978a) for the Java Sea and Lee (1974) for Hong Kong. See Sections 2 and 6.2 for global and broader references.

6.2.4 Threadfin breams, Nemipteridae (Figure 13)

Three genera have been reported from the region, namely Nemipterus, Scolopsis and Parascolopsis. The first genus is the most important in terms of quantity and commercial value. The total number of species occurring in each of the genera are 20, 11 and 2 respectively. The important species of the genus Nemipterus include N. hexodon, N. japonicus, N. marginatus, N. nematophorus, N. peronii and N. nemurus. Besides these, Scolopsis taeniopterus is also common in the commercial catch. Threadfin breams are most abundant at the intermediate depth zone ranging from 30 to 60 m (Figure 13).

It is to be noted that 10 species of this group have been studied so far, which is a relatively larger number in comparison with the other demersal fish groups studied. However, detailed research has been limited to the Gulf of Thailand (Table 4).

1/ Recently classified mostly as Paracaesio spp. and partly as Pinjalo spp.[4/]

2/ Recently classified as Symphorus spp.[4/]

3/ Recently classified as Lutjanus lutjanus[4/]

4/ Reference FAO Species Catalogue on snappers of the world. FAO Fish.Synop., (125)Vol.6:208 (Allen, 1985)

The following literature refers specifically to threadfin breams in the region: Boonwanich (1980, 1981) for the Andaman Sea, Amornchairojkul et al. (1982), Isarankura and Pariyanond (1963), Rugvichai (1977), Thumnyom and Charnprasertporn (1977), Vadhanakul (1980, 1981) for the Gulf of Thailand, Weber et al. (1977) for east Malaysian waters, Pauly and Martosubroto (1980) for the western Kalimantan waters and Arce (1981) for the Visayan Sea. See Sections 2 and 6.2 for global and broader references.

6.2.5 Bigeye snappers, Priacanthidae (Figure 14)

There is only one genus in this group comprising three species, two of which are of high commercial value: Priacanthus tayenus and P. macracanthus. The first species is more abundant than the second. Bigeye snappers are distributed widely over the region, but large concentrations are usually found at the intermediate depth zone ranging from 30 to 60 m. They are occasionally found in large concentration in deeper zones, but only in certain areas (Figure 14).

Many biological features of this group have been relatively well studied. It is notable that information for P. tayenus is available mostly in the Gulf of Thailand while that for P. macracanthus is only in the Java Sea (Table 4), a situation that needs future clarification of the reasons involved, particularly regarding their distribution patterns in the entire region.

The following literature refers specifically to bigeye snappers in the region: Chomjurai (1970) and Wetchagarun (1971) for the Gulf of Thailand and Nugroho et al. (1983) for the Java Sea. See Sections 2 and 6.2 for global and broader references.

6.2.6 Groupers, Serranidae (Figure 15)

This group is represented by nine genera and 42 species in the region. The genus Epinephelus is considered to be the most important in terms of abundance and commercial value. There are 21 species of Epinephelus in the region, four of which - E. awoara, E. megachir, E. sexfasciatus and E. tauvina - are commonly found in the commercial catch. The distribution pattern is variable by species and appears to be rather random, lacking a consistent relationship between abundance and depth of habitat (Figure 15).

Information on the biological features of this group is very limited in terms of both the number of species examined and the localities surveyed (Table 4). It is hoped that more extensive studies will be undertaken in the near future to meet industrial requirements as the importance of these species for mariculture material is rapidly growing in many countries in the region.

The following literature refers specifically to groupers in the region: Tan (1974) and Tan and Tan (1974) for the South China Sea, and Intong (1980a) for the Gulf of Thailand. See Sections 2 and 6.2 for global and broader references.

6.2.7 Croakers, Sciaenidae (Figure 16)

Fifteen genera of croakers have been reported in this region. However, the fish of commercial value are limited to a few genera and species. They are composed mostly of Johnieops sina, J. dussumieri, Otolithes ruber[1/] and Pennahia macrocephalus (Table 1) and are generally abundant in shallow waters down to 30 m deep (Figure 16).

Information on the biological features of the fish is very limited. This is due mostly to their lesser commercial value although substantial amounts are usually being landed. The fish in the Gulf of Thailand have only been intensively studied for various biological items (Table 4).

Only Phettongkam and Thasananukulkuj (1972) refer specifically to croakers in the region. See Sections 2 and 6.2 for global and broader references.

6.2.8 Lizardfish, Synodontidae (Figure 17)

Lizardfish in the region is represented by only one genus, Saurida. Among nine species found in region, the two are of high commercial value: S. wanieso[2/] and S. undosquamis. They are distributed widely in both shallow and deep waters, but the largest concentration is usually formed around the intermediate depth zone at about 30-60 m (Figure 17).

1/ A synonym, Otolithes argenteus

2/ A synonym, Saurida tumbil

Information on biological features of this group has been collected from relatively many areas in the region. However, detailed studies have only been made for the Gulf of Thailand and Manila Bay (Table 4).

The following literature refers specifically to lizardfish in the region: Boonwanich and Amornchairojkul (1982) for the Andaman Sea, Charnprasertporn (1979), Chomdej (1966), Intong (1971, 1976), Meemeskul and Boonwanich (1982) and Shinoda and Intong (1977) for the Gulf of Thailand, and Tiews, Mines and Ronquillo (1972) for the Philippines. See Sections 2 and 6.2 for global and broader references.

6.3 Shrimps[1/] (Figure 18)

The most important groups of shrimp resources in the region are Penaeus and Metapenaeus spp. It should be noted that about 70 percent of the total shrimp catch are caught in a mixture of several species. The common and commercially important species include banana prawn (Penaeus merguiensis), Indian white prawn (P. indicus), western king prawn (P. latisulcatus), giant tiger prawn (P. monodon), green tiger prawn (P. semisulcatus), Jinga shrimp, (Metapenaeus affinis), yellow shrimp (M. brevicornis), greasyback shrimp (M. ensis) and bird shrimp (M. lysianassa).

Besides these larger shrimps, small-size pelagic shrimps of the genus Acetes also occur abundantly over the entire region. They prefer the neritic environment and form dense shoals in the waters close to shore in specific areas and seasons. It should be noted here that the Western Central Pacific, including this particular region, is the most productive among the world oceans for pelagic shrimps. Major species traditionally harvested include northern mauxia shrimp (Acetes chinensis), Tsivakihini paste shrimp (A. erythraeus), Jawla paste shrimp (A. indicus), Taiwan mauxia shrimp (A. intermedius), akiami paste shrimp (A. japonicus), Alamang shrimp (A. sibogae) and Jembret shrimp (A. vulgaris). The global reviews on the biology and ecology of pelagic shrimps and the fisheries have been well documented by Omori (1974, 1975). However, too little attention has been paid so far by local institutes, and improvement is awaited as pelagic shrimps are very important not only for direct human consumption but also as food items in the fish communities in the region.

The shrimp resources as a whole are widely distributed throughout the region, but large concentrations are confined mostly to the nearshore waters especially where there is substantial discharge of fresh water (Figure 18).

Many research institutes in the region recently initiated intensive studies on the biology of commercially important species. These studies have focused mostly on several productive areas such as the Gulf of Thailand, Manila Bay, the Arafura Sea, the Java Sea and the waters along Brunei and East Malaysia. The information on biological features of various species acquired so far are given in Table 5.

In addition to the global, references mentioned in Section 2 of this paper, the following literature refers specifically to shrimps in the region.

(i) Overview

(1) Resources and fisheries:

Mistakidis (1973) and SCSP (1980) for the South China Sea, Kungvankij et al. (1973a) for the Andaman Sea, Vibhasiri (1984) for the Gulf of Thailand, Weng (1978) for Malaysia, Simpson and Chin (1978) for Sabah, Malaysia and Naamin (1984) for the Arafura Sea.

(2) Biology in general:

Holthuis (1980) for the entire region, Chaitiamvong (1980) and Vibhasiri and Kongmuang (1976, 1976a) for the Gulf of Thailand, Dwiponggo and Pauly (in press) for Indonesia, Currie (1982) for the Burnei estuary, and Tiews, Bravo and Ronquillo (1972) for Manila and San Miguel Bays.

(3) Spawning and larvae:

Kungvankij (1973) for the Andaman Sea, Pathansali et al. (1974) for the Malacca Strait, Thubtimsang (1977), Thubtimsang and Meechukhan (1976) and Thubtimsaeng (1978, 1978a) for the Gulf of Thailand and Martosubroto (1977) for the Java Sea.

1/ The English and scientific names employed in this sub-section are those defined in FAO species catalogue on shrimps and prawns of the world (Holthuis, 1980)

(ii) Specific species groups

(1) Penaeus spp.

Chaitiamvong and Thaiprayoon (1978) and Hutasing (1974) for the Gulf of Thailand, Badrudin (1978) for northwest Sumatra, Martosubroto (1977) for northwest Java, Adisusilo (1983) for Cilicap, Indonesia, and Sumiono (1983) and Naamin (1984) for Irian Jaya, Indonesia.

(2) Metapenaeus spp.

Chaitiamvong and Thaiprayoon (1977, 1978a, 1979) and Chaitiamvong and Phunwichian (1980) for the Gulf of Thailand.

(3) Acetes spp.

Omori (1974, 1975) for the entire region and Chaitiamvong (1980a) for the Gulf of Thailand.

6.4 Cephalopods[1/] (Figures 19 and 20)

The important cephalopod resources in the region are squids and cuttlefishes. Less attention has been given to the utilization of octopuses either by fishermen or consumers. Among these groups, four species of squids (Indian squid, Loligo duvauceli; swordtip squid, L. edulis; mitre squid, L. chinensis; and bigfin reef squid, Sepioteuthis lessoniana) and two species of cuttlefish (pharaoh cuttlefish, Sepia pharaonis; and curvespine cuttlefish, S. recurvirostra) are commonly found in commercial catches. They are widely distributed in both coastal and offshore waters (Figures 19-20).

Although squids and cuttlefish are generally considered to have acquired both the pelagic and demersal life forms, they are mainly caught by trawl in the region, which means that the pelagic form has been scarcely utilized. The recent development of dipnet and mid-water trap squid fishing in the Gulf of Thailand is to be noted in this regard, as it uses the pelagic form of squids in offshore waters.

Information on the biology of these resources in the region is scanty and limited to the Gulf of Thailand. They are summarized in Table 6.

In addition to the global references mentioned in the Section 2 above, the following literature refers specifically to cephalopods in the region.

(1) Overview:

Chikuni (1983) and Roper, Sweeney and Nauen (1984) for the entire region, Cholatarn (1978), Supongpan (1984) and Supongpan and Kongmuag (1976, 1976a) for the Gulf of Thailand.

(2) Squids:

Chotiyaputta (1982) for the Gulf of Thailand and Latiff (1982) for Malaysia.

(3) Loligo spp.:

Rattanaanun (1977, 1980) for the Gulf of Thailand and Marzuki (1983) and Marzuki et al. (1983) for Lombok, Indonesia.

7. DISCUSSION

It is recognized that the major fishing grounds for many important resources are widely spread out in the coastal waters of the region (Figures 3 to 20). However, intensive fishing is generally concentrated in nearshore waters shallower than 60 m deep. Fishing on some stocks like coastal tunas and king mackerels extends, on the other hand, further to offshore waters. Some demersal fish in deeper waters are also exploited on coral reefs, but the deeper parts of continental shelves generally remain unused. It is thought that the concentration of fishing in many nearshore waters in the region may be the result of the small-scale nature of the local fisheries, which depend strongly on the coast or shallower sea bottom. The availability of resources decreases, as

---

1/ The names employed refer to FAO species catalogue on cephalopods of the world (Roper, Sweeney and Nauen, 1984)

a matter of course, with increases in sea depth. However, this may not be the critical limiting factor on the expansion of fishing ground further offshore. It is believed accordingly that substantial amounts of both demersal and pelagic fish would be taken even in the offshore waters in the region where no intensive fishing has been attempted so far.

The results of a few exploratory fishing trials in offshore waters carried out in the past indicate the abundant distribution of some demersal fish stocks on the deeper part of the continental shelves. They include red snappers, groupers and threadfin breams. There may be substantial demersal fish stocks in the other offshore areas where such trials have not been made. The results of larval survey also suggest the substantial distribution of pelagic fish resources in offshore waters (Vattanachai, 1974; Termvidchakorn, 1983). In this regard, further research and studies are strongly recommended, and particularly on the relationship between inshore and offshore stocks.

As was stated in the introductory part of this paper, the geographic distribution maps presented here were prepared on a species group basis combining the information on major species in each group together. Each illustration looks like a very broad distribution over the region without any disruption or separation of the stocks. However, it is known that the distribution range generally differs substantially by species in each of the species groups referred to, as is the case for those in the Gulf of Thailand (Pramokechutima, 1978). The species composition of existing resources and of commercial catches therefore vary greatly by area. It is also well known that many species in the region are composed of several different stocks (FAO/SEAFDEC, 1985). In this regard data collection should be intensified in the future to examine the range and pattern of the distribution of specific stocks.

Another constraint is the lack of a firm basis for the species identification of many species. This is due first to the tropical nature of the resources, i.e., multispecies with lesser morphometric variation, and second to the insufficiency of the research activities carried out so far in the region. Even if the fish are limited to a few commercially important species, e.g., Decapterus spp., concensus on the species identification of their scientific names has not yet been established in the region, as was discussed at the FAO/SEAFDEC Workshop on shared stocks in Southeast Asia (FAO/SEAFDEC, 1985). It is extremely difficult to improve the situation in the immediate future due to the complexity of the subject and to the necessity of difficult work. However, the importance of continuous and intensive research and study on this particular subject needs to be re-emphasized.

A lot of biological information was summarized in this paper for a large number of species (Tables 2-6). However, a gap in the knowledge still remains in terms of the coverage of both the subjects and localities in the region. Although various intensive studies were carried out for several important stocks, they concentrated generally on a few limited areas. Knowledge is still seriously lacking in many areas, especially for the central to eastern and northern parts of the region. It is recommended that more attention be paid to the biological research of the resources, and particularly of commercially important stocks. These include mackerels, round scads, sardines, carangids, coastal tunas and shrimps. Similar activities should be expanded to the resources that have not yet been intensively harvested, but which are believed to become very important in the near future. These may include anchovies, squids, cuttlefish and octopuses.

Among the many biological items, growth, mortality, feeding habit and spawning season have been relatively intensively studied. However, the data are generally limited to the western part of the region, the Gulf of Thailand, the Malacca Strait and the Java Sea. Furthermore, spawning ground, first maturity and life span, which are also very important parameters for assessment purposes, have not been intensively studied. It should be noted again that knowledge on these features in the central to eastern and northern parts of the region remains meager and the importance of future study needs to be emphasized.

It is to be noted that the magnitude and quality of the studies vary according to the area surveyed. This may be the result of many factors including the complexity of the environment and the resources in the areas surveyed, sampling techniques and the method of analysis employed, etc. It is hoped, in this regard, that knowledge on the biology of important resources will be obtained for the entire region, first on a well-balanced basis for geographic coverage and second on a compatible basis in terms of the minuteness and preciseness among the areas/institutes concerned. It is expected that this paper will be useful as a guidepost in this respect.

The information on the migratory patterns of pelagic fish is also very sparse and fragmentary. Although the available information has been compiled and given in Figure 21, it is very tentative for the reasons mentioned above. Through tagging experiments carried out by the research institute in Thailand, the Indo-Pacific mackerel (Rastrelliger brachysoma) was confirmed to migrate seasonally in the Gulf of Thailand. The experiments also suggest that a fairly demarcated range of the distribution of the fish exists in the Gulf, from the inner part to the northwest coast, which

falls entirely in Thai jurisdictional waters. The fish in the Malacca Strait, on the other hand, migrate seasonally across the jurisdictional border between Thailand and Malaysia. There are many stocks in the other waters of the region that show short-distance migration and displacement within local waters in conjunction with spawning and feeding. These include anchovies in Manila Bay and the Strait of Singapore; sardines in Manila Bay, the Samar Sea and the Bali Strait; many demersal fish in the waters off Sarawak and Peninsular Malaysia and the northwestern part of the South China Sea. Shrimps in the Brunei Bay show a similar displacement pattern, and the same is probably true for the many other shrimp stocks in the region.

The distribution and migration patterns of fish sometimes raise further complicated problems in harvesting and managing these resources as shared stocks in both the migratory and non-migratory as well as the "transboundary" modes. This has become one of the focal points of resource management for all the countries in the region since national jurisdiction was extended to 200 miles. In fact, there are probably at least 40 stocks currently being shared by two or more countries as identified at the FAO/SEAFDEC Workshop on shared stocks in Southeast Asia (FAO/SEAFDEC, 1985). The biological aspects of these stocks were discussed and the major direction of future research was clarified at the Workshop. The principal requirements for the research of shared stocks are quite identical to those for stock assessment in general. However, the priority areas specifically identified were the geographic limits of distribution, migration and displacement and the location and timing of spawning. Improvement in the quality and timeliness of fishery statistics is another essential issue to be tackled in the immediate future, as was also discussed at the Workshop. It is hoped, in this regard too, that this paper will stimulate an improvement in research in the region.

The strengthening of international cooperation among the institutes and the scientists concerned should also be pursued. Collaboration in research and study, including frequent and close communication, would undoubtedly push forward not only resources appraisal and management but also marine sciences in general for the region.

Finally, apart from the intra-species biological features of important fish resources mentioned above, the importance of inter-species functions in the fish community, including those of trash fish, also needs to be emphasized. In the combination of multispecies resources and multigear fisheries, both distinctive features of this particular region, as was already mentioned, the species composition of the fish community in the region could easily be considerably changed. In fact, there have been several clear indications of this phenomenon in the demersal fish community in the Gulf of Thailand (Ritragsa, 1976; Pauly, 1979). It has generally been accepted by many scientists that the decline in the commercially important larger size finfish would have resulted in a substantial increase in the other species, e.g., cephalopods and smaller size trash fish. However, studies on these inter-species aspects of the fish community have scarcely been made so far due first to the lack of adequate information and second to the extremely high complexity of the subject and approach (Pauly and Murphy (eds.), 1982). This is a long-term target, and the scientists and institutes in the region need to keep this in mind for the future research.

8. REFERENCES

Adisusilo, S., 1983 — A review on size maturity and spawning season of banana prawn (Penaeus merguiensis de Man) at Cilacap and adjacent waters. Mar.Fish.Res.Rep.,Jakarta, (29):97-103 (in Indonesian)

Aglen, A., et al., 1981 — Surveys of the marine fish resources of Peninsular Malaysia, June-July 1980. Report on Surveys with the R/V Dr Fridtjof Nansen. Bergen, Institute of Marine Research, 70 p.

Allen, G.R., 1985 — FAO species catalogue. Vol. 6. Snappers of the world. An annotated and illustrated catalogue of lutjanid species known to date. FAO Fish.Synop., (125)Vol.6:208 p.

Amaragsa, C., 1986 — Size composition of Leiognathus bindus (Valenciennes), L. leuciscus (Tunther), and Pentaprion longimanus (Cantor). Paper presented to IPFC Conference, Australia, 1968 (mimeo)

Amornchairojkul, S. and T. Boonwanich, 1982 — Growth and mortality of Nemipterus japonicus off the west coast of the Gulf of Thailand, (Chumporn-Ko Samui), 1981. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (7/2525):12 p. (in Thai)

Aoyama, T., 1973 — The demersal fish stocks and fisheries of the South China Sea. Rome, FAO, South China Sea Fisheries Development and Coordinating Programme, SCS/DEV/73/3:80 p.

Arce, F.M., 1981 Distribution and relative abundance of Nemipterids and Carangids (Pisces: Nemipteridae and Carangidae) caught by trawl in Visayan Sea with notes on the biology of Nemipterus oveni and Selaroides leptolepis. Master Thesis submitted to the College of Arts and Sciences, University of the Philippines (Unpubl.)

Atmadja, S.B., 1982 Gonad indices and sex ratio of round scad (Decapterus maruadsi) in the Arafura Sea. Mar.Fish.Res.Rep.,Jakarta, (24):37-43 (in Indonesian)

Badrudin, M., 1978 Preliminary study on the age, growth, gonad maturation of tiger shrimp, Penaeus monodon Fabr. in Meulabok waters and off west coast of Aceh. In Proceedings of the Second Seminar on shrimp fisheries, Jahark, March 1977, pp. 135-44 (in Indonesian)

_____, 1978a Stock of goat fish, Upensus sulphureus in the Java Sea with notes on some biological aspects. In Proceedings of the Symposium on the modernization of small scale fisheries. Jakarta, 27-30 June, 1978, pp. 1-7 (in Indonesian)

Beales, R.W., et al., 1982 Investigations into fisheries resources in Brunei. Monogr.Brunei Mus.J., (5):1-204

Bhatia, U. and V. Boonragsa, 1980 Biological statistics of small tuna along the west coast of Thailand. Phuket, Phuket Marine Fisheries Station, 20 p. (MS report, in Thai)

Bhatia, U., et al., 1979 Marine fisheries in the Andaman Sea. In Report on the results of Seminar on the Marine Fisheries Development Project, edited by V. Hongskul. Tech.Rep.Mar.Fish. Div.,Bangkok, (79/1):230 p. (in Thai)

Blanc, M., 1960 Mission hydrobiologique et océanographique au Cambodge (février-mars 1959). Cah. Pacif.,Paris, (2):33-62

Boonprakob, U., 1965 The identification of pelagic eggs and larvae of chub mackerel, Rastrelliger spp., in the Gulf of Thailand with additional study on their distribution. Contrib.Mar.Fish. Lab.,Bangkok, (4):115-51 (in Thai)

_____, 1967 Study on the fecundity of the Indo-Pacific mackerel, Rastrelliger spp., in the Gulf of Thailand. Proc.IPFC, 12(2):124-38

_____, 1972 Conservation measures for Pla-tu, Rastrelliger neglectus (van Kampen) of the Gulf of Thailand. Thai Fish.Gaz., 25(3):303-44 (in Thai, with English abstr.)

Boonragsa, V., 1984 Preliminary study on the present status of pelagic fisheries in the Andaman Sea. Paper presented to the Marine Fisheries Seminar, Bangkok, 4-7 September 1984, 46 p. (in Thai)(mimeo)

Boonragsa, V., et al., 1984 Biological studies and the present status of pelagic fisheries resources in the Andaman Sea. Paper presented to the Marine Fisheries Seminar, Bangkok, 4-7 September 1984 (mimeo)

Boonwanich, T., 1980 Length composition, growth and sex ratio of Nemipterus mesoprion in the Gulf of Thailand (January 1978 - March 1979). Tech.Rep.Demersal Fish.Unit Mar.Fish.Div., Bangkok, (4/1980)

_____, 1981 Size composition and growth of threadfin bream (Nemipterus nematophorus) in the Gulf of Thailand, January 1978 - March 1979. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div., Bangkok, (3/1981)

Boonwanich, T. and S. Amornchairojkul, 1982 Length composition of Saurida undosquamis and S. elongata in the west coast of the Gulf of Thailand, 1981 (Chumphon-Koh Samui). Tech.Rep. Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (13/1982):16 p. (in Thai)

Burma, Socialist Republic of the Union of, 1982 Notes on fisheries in Burma. Rangoon, Ministry of Agriculture and Forests, January 1982

Bussarawitch, N., 1984 Some biological aspects of the Indian mackerel (Rastrelliger kanagurta) in the Indian Ocean coast of Thailand. Technical report. Phuket, Phuket Marine Fisheries Station

_____, 1984a Fecundity of the Indo-Pacific mackerel, Rastrelliger brachysoma along the west coast of Thailand. Technical Report. Phuket, Phuket Marine Fisheries Station, 15 p. (in Thai)

Bussarawitch, N., 1984b Biological aspects of Rastrelliger brachysoma on the Indian Ocean coast of Thailand. Technical report. Phuket, Phuket Marine Fisheries Station, 25 p. (in Thai)

Chaitiamvong, S., 1980 The biology of the penaeid shrimps of Thailand. In Report of the Workshop on the biology and resources of penaeid shrimps in the South China Sea area. Part 1. 30 June - 5 July 1980, at Kota Kinabalu, Sabah, Malaysia. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/80/26:93-122

_____, 1980a Biological studies on planktonic shrimp and shrimp-like in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1980):18 p. (in Thai)

Chaitiamvong, S. and S. Phunwichian, 1980 Biological studies on Metapenaeus intermedius Kishinouye in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1980):23 p. (in Thai)

Chaitiamvong, S. and S. Thaiprayoon, 1977 Biological study on Metapenaeus mutatus Lanchester, in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1977):23 p. (in Thai)

_____, 1978 Biological study on Penaeus semisulcatus de Haan, in the Gulf of Thailand. Annu. Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1978):30 p. (in Thai)

_____, 1978a Biological studies on Metapenaeus mutatus Lanchester, in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1978):23 p. (in Thai)

_____, 1979 Biological study on Metapenaeus ensis (de Haan) in the Gulf of Thailand. Annu.Rep. Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1979):23 p. (in Thai)

Chantarasri, D., 1980 A study on age and growth of the round scad (Decapterus maruadsi Temminck et Schlegel) caught in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div., Bangkok, (15):48 p. (in Thai)

Chantawong, T., et al., 1984 Trawl fisheries and the present status of demersal resources in the Andaman Sea. Paper presented to the Marine Fisheries Seminar, Bangkok, 4-7 September 1984 (mimeo)

Charnprasertporn, T., 1979 Length-weight relationship of lizard fish (Saurida elongata) in the Gulf of Thailand. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (5/2522):13 p. (in Thai)

Chayakul, R., et al., 1983 Abundance of fish larvae in the west coast of the Gulf of Thailand. Tech. Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (31):23 p. (in Thai)

Chee, P.E., 1980 The reproductive pattern of Rastrelliger kanagurta. In Proceedings of the Second Annual Seminar. Kuala Lumpur, Malaysian Society of Marine Science, pp. 33-44

Cheunpan, A., 1981 Fecundity of round scad (D. maruadsi) in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (25):12 p. (in Thai)

_____, 1984 Sexual maturity, size at first maturity and spawning season of longtail tuna (T. tonggol), eastern little tuna (E. affinis) and frigate mackerel (A. thazard) in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Sub-Div.Mar.Fish.Div.,Bangkok, (43):22 p. (in Thai)

Chiampreecha, B., 1978 Biological studies on tuna and tuna-like fishes in the west coast of the Gulf of Thailand and off east coast of Peninsular Malaysia. Tech.Rep.Pelagic Fish.Unit Mar. Fish.Div.,Bangkok, (4):25 p.

Chikuni, S., 1983 Cephalopod resources in the Indo-Pacific region. FAO Fish.Tech.Pap., (231):264-305

Cholatarn, C., 1978 Study on distribution and abundance of some species of cephalopods in the Gulf of Thailand. Tech.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (26/1978)

Chomdej, V., 1966 Size composition, sex ratio and spawning periods of greater lizard fish, Saurida tumbil, in the Gulf of Thailand. Contrib.Mar.Fish.Lab.,Bangkok, (5):21 p.

Chomjurai, W., 1970 Some aspects on the biology of spot-finned bigeye (Priacanthus tayenus) in the inner part of the Gulf of Thailand (1968-1969). Paper presented to the Symposium on marine fisheries, Bangkok, 26-27 January 1970. Bangkok, Marine Fisheries Laboratory, 15 p. (mimeo)

Chong, B.J. and C.W. Chua, 1974 Growth, age determination and spawning of the ikan kembun, Rastrelliger neglectus (van Kampen) in the northern Strait of Malacca. Malay.Agric.J., 49:344-56

Chotiyaputta, C., 1982 Squid fishery of Thailand. FAO Fish.Rep., (275):124-34

Chua, T.E. and J.A. Mathias (eds), 1975 Coastal resources of West Sabah. Penang, Malaysia, University Sains Malaysia

Chullasorn, S., 1979 Mackerel resources in the Gulf of Thailand. Thai Fish.Gaz., 32(1):55-69 (in Thai)

_____, 1979a Sardines. In Results of the Seminar on Marine Fisheries Development Program. Bangkok, Marine Fisheries Division, Department of Fisheries, Annex 4-6, 3 p. (in Thai)

Chullasorn, S. and S. Yusukswad, 1977 Preliminary report on the fisheries biology of the round scad (Decapterus spp.) in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div., Bangkok, (2):34 p.

Chullasorn, S., K. Chotiyaputta and R. Chayakul, 1973 Preliminary studies on the narrow-barred Spanish mackerel, Scomberomorus commerson (Lacépède, 1802) in the Gulf of Thailand. Tech.Rep. Pelagic Fish.Unit Fish.Div.,Bangkok, (1973):47 p. (in Thai)

Chung, R.T. and M.T.L. Chiu, 1980 The purse seine fishery of Hong Kong based on an investigation conducted in 1975-1976. Proc.IPFC, 19:437-60

Collette, B.B. and C.E. Nauen, 1983 FAO species catalogue. Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. FAO Fish.Synop., (125)Vol.2:137 p.

Currie, D., 1982 Studies on the biology of penaeid prawns and the prawn fishery in the Brunei estuary. In Investigations into fisheries resources in Brunei. Monogr.Brunei Mus.,J., (5): 134-203

Departemen Pertanian, Indonesia, 1983 Estimates of marine resource potential of the Indonesian waters. Jakarta, Department of Agriculture, Indonesia, 25 p. (in Indonesian)

Department of Fisheries, Thailand, 1984 Purse seine fisheries and the present status of pelagic fish resources in the Gulf of Thailand. Paper presented to the Marine Fisheries Seminar, Bangkok, 4-7 September 1984, 68 p. (mimeo in Thai)

Dhebtaranon, P., 1973 Length frequency distribution of Stolephorus heterolobus (Rüppel) in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (1973):15 p.

Dhebtaranon, Y. and K. Chotiyaputta, 1974 Review of the mackerel fishery (Rastrelliger spp.) in the Gulf of Thailand. Proc.IPFC, 15(3):265-86

Druzhinin, A.D., 1968 Indian mackerel, Rastrelliger spp. in Burmese waters. Proc.IPFC, 13(2):59-81

Druzhinin, A.D. and U.P. Hliang, 1972 Observation on the trawl fishery of southern Burma. Proc.IPFC, 13(3):151-209

Dwiponggo, A., 1974 The fishery for and preliminary study on the growth rate of "Lemuru" (oil sardine) at Muntjar, Bali Strait. Proc.IPFC, 15(3):221-40

Dwiponggo, A. and D. Pauly, An atlas of the growth, mortality and recruitment of Indonesian fishes. ICLARM Tech.Rep. (in press)

Dwiponggo, A. and J.C.B. Uktolseya, 1974 A lemeru, Sardinella longiceps, survey around the western part of the Little Sunda Island. Proc.IPFC, 15(3):211-20

FAO, 1975 Offshore fishery development project. Republic of Viet Nam. An analysis of data of the purse seine and sonar survey 1971-1972, with estimates of the abundance and potential yield of Vietnamese offshore pelagic resources. Report prepared for the Government of Viet Nam by FAO of the UN acting as executing agency for the UNDP. Based on the work of P. Gonzales-Alberdi and C. Berude. Rome, FAO/UNDP, FI:SF/VIE3, Technical report 5:39 p.

_____, 1980 Fishery country profile. Democratic Kampuchea. Rome, FAO, FID/CP/KHM Rev.1:4 p.

FAO/SEAFDEC, 1985 Report of the FAO/SEAFDEC Workshop on shared stocks in Southeast Asia. Bangkok, 18-22 February 1985. FAO Fish.Rep., (337):97 p.

FAO/UNDP, 1985 Marine Fishery Resources Management in the Bay of Bengal Project, Mackerels in the Malacca Strait. Colombo, Sri Lanka, Bay of Bengal Programme RAS/81/051, BOBP/WP/30: 63 p.

Fisheries Research Institute, Malaysia, 1984 Annual report for 1982. Annu.Rep.Fish.Res.Inst.Glugor, Penang, (1982)

Gafa, B., 1982 Some biological aspects of mackerel (Rastrelliger kanagurta) caught in the Makassar Strait during early monsoon periods. Mar.Fish.Res.Rep.,Jakarta, (23):91-101 (in Indonesian)

Gafa, B. and B. Sadhotomo, 1983 Investigation on growth and some biological parameters of John snapper (Lutjanus johni) in the South Kalimantan waters. Mar.Fish.Res.Rep.,Jakarta, (29):89-96 (in Indonesian)

Holthuis, L.B., 1980 FAO species catalogue. Vol. 1. Shrimps and prawns of the world. An annotated and illustrated catalogue of species of interest to fisheries. FAO Fish.Synop., (125):Vol.1:261 p.

Hutasing, R., 1974 Study on the fecundity of white shrimp, Penaeus merguiensis de Man. Annu.Rep.Rayong Mar.Fish.Stn.Rayong,Thailand, (1974):162-82 (in Thai)

Ingles, J. and D. Pauly, 1984 An atlas of the growth, mortality and recruitment of Philippine fishes. ICLARM Tech.Pap., (13):127 p.

Intong, S., 1971 Biology of lizard fish, Saurida undosquamis (Richardson) in the inner Gulf of Thailand. Paper presented to the Second Symposium on marine fisheries, Bangkok, Marine Fisheries Laboratory, 19-20 April 1971, 27 p. (mimeo)

_____, 1976 Foods and feeding of lizard fish, Saurida undosquamis (Richardson) off Cholburi in the inner Gulf of Thailand, October 1971 - August 1972. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (4/2519):26 p. (in Thai)

_____, 1980 Food composition in the stomach of malabar snapper, Lutjanus malabaricus, in the Gulf of Thailand, 1978. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (10/1980)

_____, 1980a Food composition in the stomach of sea bass grouper, Epinephelus sexfasciatus, in the Gulf of Thailand, 1978. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (11/1980):22 p. (in Thai)

Isara, P., 1972 Length frequency distribution of the Stolepherus heterolobus (Rüpell) in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (1968-1972). Part 1:265-88

Isarankura, A.P., 1968 Present status of demersal fish resources in the Gulf of Thailand from the analysis of experimental trawl survey. Tech.Rep.Demersal Fish.Unit Dep.Fish.,Bangkok, 20 p. (unpublished, in Thai)

Isarankura, A.P. and S. Pariyanond, 1963 Preliminary studies on the life history of threadfin bream, Nemipterus hexodon, in the Gulf of Thailand. Contrib.Dep.Fish.,Bangkok, (1):28 p.

Kangvankij, P., 1970 Biological study of red snapper, Lutjanus sanguineus. Report submitted to Southeast Asian Fisheries Development Center (SEAFDEC), Singapore, September 1970 (mimeo)

Khin, U., 1948 Fisheries in Burma. Government of the Union of Burma, Rangoon, Burma, Superintendent, Govt. Printing and Stationery, 121 p.

Kikawa, S. and M.G. Ferraro, 1967 Maturation and spawning of tunas in the Indian Ocean. Proc.IPFC, 12(2):65-78

Klinmuang, H., 1978 Preliminary studies on the biology of tunas in the west of the Gulf of Thailand and off the east coast of Peninsular Malaysia. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div., Bangkok, (1978):27 p.

_____, 1979 Pla-O. In Results of the Seminar on Fisheries Development Program organized by Marine Fisheries Division, Department of Fisheries, Bangkok, Annex 4.8, 4 p. (in Thai)

_____, 1981 Length frequency distribution and length-weight relationship of tunas in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok (24):33 p.

Kume, S., 1973 Tunas resources in the South China Sea. Rome, FAO, South China Sea Fisheries Development and Coordinating Programme, SCS/DEV/73/4:18 p.

Kungvankij, P., et al., 1973 Observation on the spawning season of three economically important species of shrimps from the Indian Ocean coast of Thailand, estimated from gonad index. Contrib.Phuket Mar.Fish.Stn.,Phuket, (4):8 p.

_____, 1973a A survey of the distribution and abundance of economically important shrimp along the Indian Ocean coast of Thailand. Contrib.Phuket Mar.Fish.Stn.,Phuket, (3):9 p. (in Thai)

Kyokuyo Co., Ltd., 1972 Trawling survey in the South China Sea and the Gulf of Thailand (1969-1970-1971). Executed for the Government of the Republic of Vietnam. FI:SF/VIE 3 Final report. Tokyo, Kyokuyo Co., Ltd., 236 p.

Lai-shing, A., 1968 Status of the Hong Kong trawling industry. Proc.IPFC,13(3):270-97

Latiff, M.S.B.S.A., 1982 Malaysian squid resources. FAO Fish.Rep., (275):118-20

Latiff, M.S.B.S.A., et al., 1974 Demersal fish resources in Malaysian waters. 2. Trawl survey of the coastal waters off the west coast of Peninsular Malaysia, 12 December 1970 to 22 January 1971. Fish.Bull.Minist.Agric.Malays., (3):41 p.

Lee, C.K.C., 1974 The reproduction, growth and survival of Upeneus moluccensis (Bleeker), in relation to the commercial fisheries in Hong Kong. Hong Kong Fish.Bull., (4):17-32

Magnusson, J., 1970 Report on assignment as marine biologist with the UNDP(SF)/FAO Deep-Sea Fishing Development Project in the Philippines (January 1966 - June 1969). Rome, FAO, UNDP/SF Project PHI/11:90 p.

Marr, J.C., 1976 Fishery and resource management in Southeast Asia. Washington, D.C., Resources for the Future, Programme of International Studies of Fishery Arrangements. RFF/PISFA Pap., (7):62 p.

Martosubroto, P., 1977 Spawning and growth of banana shrimp, Penaeus merguiensis de Man and endeavour shrimp, Metapenaeus ensis de Haan in the waters off Krawang Peninsula (West Java). In Proceedings of the Second Seminar on shrimp fisheries, Jakarta, march 1977, pp. 7-21 (in Indonesian)

_____, 1982 Fishery dynamics of the demersal resources of the Java Sea. Ph.D. Thesis. Dalhousie University, Halifax, Canada, 238 p. (Unpubl.)

Marzuki, S., 1983 Some biological aspects on squids, Loligo spp. in Lombok (West Nusa Tenggara). Mar. Fish.Res.Rep.,Jakarta, (17):25-37 (in Indonesian)

Marzuki, S.,et al., 1983 Investigation on the biological aspects of squid (Loligo spp.) in Lombok, West Nusa Tenggara. Mar.Fish.Res.Rep.,Jakarta, (27):25-37 (in Indonesian)

Meemeskul, Y., 1978 Relationship between total length, body length, girth length and girth factors of some demersal fishes in the Gulf of Thailand, 1978. Tech.Rep.Demersal Fish.-Unit Mar. Fish.Div.,Bangkok, (2/1978):10 p. (in Thai)

Meemeskul, Y., 1982 Species and size composition of trash fish from otter board trawl monitoring survey by M.V. Pramong 2 and Pramong 9 in the Gulf of Thailand, 1980. Tech.Rep.Demersal Fish. Unit Mar.Fish.Div.,Bangkok, (5/1982):34 p. (in Thai)

Meemeskul, Y. and T. Boonwanich, 1982 Length, growth and mortality of lizard fish, Saurida undosquamis, in the Gulf of Thailand, 1979-1980. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (6/1982):25 p. (in Thai)

Menasveta, D., 1980 Resources and fisheries of the Gulf of Thailand. Samuutprakan, Thailand, SEAFDEC Training Department, (TD/TRB8):104 p.

Menasveta, D., S. Shindo and S. Chullasorn, 1973 Pelagic fishery resources of the South China Sea and prospects for their development. Rome, FAO, South China Sea Fisheries Development and Coordinating Programme, SCS/DEV/73/6:68 p.

Mistakidis, M.N., 1973 The crustacean resources and related fisheries in the countries bordering the South China Sea. Rome, FAO, South China Sea Fisheries Development and Coordinating Programme, SCS/DEV/73/7:39 p.

Morgan, J.R. and M.J. Valencia, 1983 Atlas for marine policy in Southeast Asian seas. Berkeley, California, University of California Press, 144 p.

Naamin, N., 1984 Shrimp fisheries and population dynamics of banana shrimp, Penaeus merguiensis de Man, in the Arafura Sea. Ph.D. Thesis. Bogor Agricultural University, Bogor, Indonesia, 324 p. (in Indonesian, unpubl.)

Naiyanetr, P., 1963 Preliminary studies on life history of pampano (Caranx leptolepis) in the Gulf of Thailand. Contrib.Dep.Fish.,Bangkok, (2):19 p. (in Thai)

Nguyen, T.N., 1972 The size composition and length-weight relationship of commercial demersal fishes in the South China Sea. Graduation thesis submitted to the Marine Fisheries Research Department, SEAFDEC, Singapore, 25 p.

Nugroho, D., et al., 1983 Studies on the growth and mortality of the bigeye snapper, Priacanthus macracanthus, of the north coast of Java. Mar.Fish.Res.Rep.,Jakarta, (27):9-11 (in Indonesian)

Nupetch, L., 1984 Length frequency distribution, age and growth determination of yellow striped trevally, Caranx leptolepis caught in the Gulf of Thailand. Tech.Rep.Pelagic Fish. Unit Mar.Fish.Div.,Bangkok, (47):21 p. (in Thai)

Nupetch, L., et al., 1983 The age, growth determination of hardtail scad, Megalaspis cordyla, caught in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (38):20 p. (in Thai)

Omori, M., 1974 The biology of pelagic shrimps in the ocean. Adv.Mar.Biol., 12:233-324

_____, 1975 The systematics, biogeography and fishery of epipelagic shrimps of the genus Acetes (Crustacea, Decapoda, Sergestidae)

Ordonez, J.A., et al., 1973 Zooplankton distribution off Mindoro Island and Balayan Bay, Luzon island, Philippines-South China Sea. Philipp.J.Fish., 11(1-2):23-35

Pathansali, D., 1961 A preliminary report on the Rastrelliger fishery in Malaysia. Proc.IPFC, 2(2): 37-45

_____, 1967 Observations on the gonad maturity stages of female Rastrelliger kanagurta Cuvier. Proc.IPFC, 12(2):112-5

Pathansali, D., et al., 1974 Demersal fish resources in Malaysian waters. Trawl survey of the coastal waters off the east coast of West Malaysia. Fish.Bull.Minist.Agric.Fish.Malays., (1): 46 p.

Pauly, D., 1979 Theory and management of tropical multispecies stocks; a review on the Southeast Asian demersal fisheries. ICLARM Stud.Rev., (1):1-35

Pauly, D. and P. Martosubroto, 1980 The population dynamics of Nemipterus marginatus (Cuvier & Val.) off western Kalimantan, South China Sea. J.Fish.Biol., 17:263-73

Pauly, D. and G.I. Murphy (eds.), Theory and management of tropical fisheries. Proceedings of the 1982 ICLARM/CSIRO Workshop on the theory and management of tropical multispecies stocks, Cronulla, Australia, 12-21 January 1981. ICLARM Conf.Proc., (9):360 p.

Phettongkam, M. and W. Thasananukulkuj, On the biology of Sciaena russelli (Cuvier et 1972 Valenciennes) in the inner Gulf of Thailand. Paper presented at the Third Symposium on marine fisheries, Bangkok, Marine Fisheries Laboratory, 18 p. (mimeo)

Piyathirathitivorakul, P., A study on age, growth and length frequency distribution of the one 1983 finlet scad (Caranx mate Cuvier) caught in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (30):21 p. (in Thai)

_____, Age, growth and length frequency distribution of the round scad (Decapterus maruadsi 1983a Temminck et Schlegel) in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar.Fish. Div.,Bangkok, (32):27 p. (in Thai)

Pong, L.Y., Demersal fish resources in Malaysian waters. 14. Fourth trawl survey off the west 1981 coast of Peninsular Malaysia (17 October - 18 November 1978). Fish.Bull.Minist.Agric. Malays., (26):20 p.

Pong, L.Y., et al., Demersal fish resources in Malaysian waters. 12. Third trawl survey off the 1976 coast of Sarawak (16 July - 8 August 1975). Fish.Bull.Minist.Agric.Malays., (14):35 p.

Pramokechutima, S., Distribution and abundance of some demersal fishes in the Gulf of Thailand, 1978 1977. Tech.Rep.Demersal Fish.-Unit Mar.Fish.Div.,Bangkok, (3/1978)

Rattanaanun, T., Biological studies on Loligo duvaucelii d'Orbigny in the Gulf of Thailand. Tech. 1977 Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (10/2520):10 p. (in Thai)

_____, Biological studies on Loligo duvaucelii d'Orbigny (Cephalopoda; Loliginidae) in the 1980 Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1980):8 p. (in Thai)

Ritragsa, S., Results of the studies of demersal fish resources in the Gulf of Thailand from 1976 trawling survey 1963-1972. In Proceedings of the Seminar on fisheries resources and their management in Southeast Asia, Berlin, 19 November to 6 December 1974, edited by K. Tiews. Bonn, Drükerei Heing Lammerich, pp. 156-68

Ritragsa, S., D. Dhamniyom and S. Sitthichaikasem, An analysis of demersal fish catches taken from 1968 the otterboard trawling survey in the Gulf of Thailand, 1966. Contrib.Mar.Fish.Lab., Bangkok, (11):104 p.

Ritterbush, S.W., An assessment of the population biology of the Bali Strait lemuru fishery. Mar. 1975 Fish.Res.Rep.,Jakarta, (1):1-38

Ronquillo, I.A., Synopsis of biological data on Philippine sardines (Sardinella perforata, 1960 S. fimbriata, S. sirm, S. longiceps). In Proceedings of the World Scientific Meeting on the biology of sardines and related species. Rome, 14-21 September 1959, edited by H. Rosa Jr and G. Murphy. Rome, FAO, Vol. 2, pp. 453-95

_____, A contribution to the biology of Philippine tunas. FAO Fish.Rep., (62):Vol.3:1683- 1963 752

_____, A review of the round scad fishery in the Philippines. Proc.IPFC, 15(3):351-75 1974

Roper, C.F.E., M.J. Sweeney and C.E. Nauen, FAO species catalogue. Vol. 3. Cephalopods of the 1984 world. An annotated and illustrated catalogue of species of interest to fisheries. FAO Fish.Synop., (125)Vol.3:277 p.

Rugvichai, K., Some biological aspects of threadfin bream. Tech.Rep.Demersal Fish.-Unit Mar.Fish. 1977 Div.,Bangkok, (1/1977):26 p. (in Thai)

SCSP (South China Sea Fisheries Development and Coordinating Programme), Report of the Workshop on 1976 the fishery resources of the Malacca Strait. Part 1. Jakarta, 29 March - 2 April 1970. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/76/2, Part 1:89 p.

SCSP (South China Sea Fisheries Development and Coordinating Programme), Report of the Workshop on
1976a the fishery resources of the Malacca Strait. Part 2. Jakarta, 9 March - 2 April 1976. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/76/6, Part 2:113 p.

_____, Report on the BFAR/SCSP Workshop on the fisheries resources of the Visayan and
1976b Sibuyan Sea areas. Iloilo, Philippines, 18-22 October 1976. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/76/7:26 p.

_____, Report on the BFAR/SCSP Workshop on the fishery resources of the Sulu Sea, Bohol Sea
1977 and Moro Gulf areas. Cagayan de Oro, Philippines, 25-29 April 1977. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/77/11:58 p.

_____, Report on the Workshop on the demersal resources of the Sunda Shelf. Part 1.
1977a Penang, Malaysia, 31 October - 6 November 1977. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/12:44 p.

_____, Report of the Workshop on the demersal resources of the Sunda Shelf. Part 2.
1977b Penang, Malaysia, 31 October - 6 November 1977. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/13:120 p.

_____, Pelagic resources evaluation. Report of the Workshop on the biology and resources
1978 of mackerels (Rastrelliger spp.) and round scad (Decapterus spp.) in the South China Sea. Part 1. Penang, Malaysia, 7-11 November 1977. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/78/17:46 p.

_____, Report of the Workshop on management of resources of the Sunda Shelf, Malacca Strait
1978a and related areas. Manila, Philippines, 7-9 December 1977. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/78/18:14 p.

_____, Report of the BFAR/SCSP Workshop on the fisheries resources of the Pacific coast of
1978b the Philippines. Naga City, Philippines, 18-22 September 1978. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/78/19:48 p.

_____, Report of the Workshop on demersal and pelagic fish resources of the Java Sea.
1979 Semarang, Indonesia, 5-9 December 1978. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/79/20:60 p.

_____, Report of the BFAR/SCSP Workshop on the fishery resources of the north and western
1979a coast of Luzon. Manila, Philippines, 18-20 April 1979. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/79/22:57 p.

_____, Report of the Workshop on the biology and resources of Penaeid shrimps in the South
1980 China Sea area. Part 1. Kota Kinabalu, Sabah, Malaysia, 30 June - 5 July 1980. Manila, South China Sea Fisheries Development and Coordinating Programme, SCS/GEN/80/26:162 p.

SEAFDEC (Southeast-Asian Fisheries Development Center), Report of the Seminar on stock assessment
1982 of pelagic resources with emphasis on shared stocks. Bangkok, Thailand, 10-14 August 1981, Bangkok, SEAFDEC, 219 p.

_____, Fishery statistical bulletin for the South China Sea area, 1982. Fish.Stat.Bull.
1984 South China Sea Area, (1982):268 p.

Sadhotomo, B., et al., On the estimation of growth, mortality parameter and yield per recruit for
1983 round scad (Decapterus maruadsi) in the Java Sea. Mar.Fish.Res.Rep.,Jakarta, (27):1-8 (in Indonesian)

Saeger, J., et al., First report of the Indonesian-German demersal fisheries project. Results of
1976 a trawl survey in the Surda shelf area. Contrib.Mar.Fish.Res.Inst., (1):1-46

Shindo, S., General review of the trawl fishery and the demersal fish stocks of the South China
1973 Sea. FAO Fish.Tech.Pap., (120):49 p.

_____, The marine fish resources and fisheries in Southeast Asia. FAO Fish.Tech.Pap. (in preparation)

Shindo, S. and S. Chullasorn, Economically important marine fishes in the Southeast Asian waters.
1980 SEAFDEC Text/Ref.Book Ser., (17):97 p.

Shinoda, M. and S. Intong, 1977 Size frequency distribution of lizard fish, Saurida undosquamis (Richardson) in the inner Gulf of Thailand. Bull.Jap.Soc.Sci.Fish., 44(1):1-6

Shinoda, M., P.Y. Lim and S.M. Tan, 1978 Preliminary study of trash fish landed at Kangkar fish market in Singapore. Bull.Jap.Soc.Sci.Fish., 44(6):595-600

Shinoda, M., et al., 1979 A method for estimating the best cod-end mesh size in the South China Sea area. Bull.Choshi Mar.Lab.Chiba Univ., (11):65-80

Simpson, A.C. and P.K. Chin, 1978 The prawn fisheries of Sabah, Malaysia. Fish.Bull.Minist.Agric. Malays., (22):1-25

Sitthichokpun, S., 1970 A study on the spawning period and spawning grounds of Stolephorus spp. off the western coast of the Gulf of Thailand, 1968-1969. Paper submitted to Department of Fisheries, Bangkok, 48 p. (in Thai)

Somjaiwong, D., et al., 1970 Progress report of the tagging of Indo-Pacific mackerel (R. neglectus), 1967. Rearing experiments. Paper presented to the First Seminar of Marine Fisheries Laboratory, Bangkok, 22 p. (mimeo)

Suchondhamarn, P., et al., 1970 Estimation of age and growth of chub mackerel, Rastrelliger neglectus (van Kampen) in the western Gulf of Thailand. In The Kuroshio: a Symposium on the Japan Current, edited by J.C. Marr. Honolulu, East-West Center Press, pp. 471-7

Sudradjat, A. and D. Nugroho, 1983 Study on the biology of yellow striped trevally, Selaroides leptolepis (Cuv. and Vab.) in the Sunda Shelf waters. Mar.Fish.Res.Rep.,Jakarta, (29): 79-96 (in Indonesian)

Sujastani, T., 1974 Population dynamics of mackerels in the Java Sea. Mar.Fish.Res.Rep.,Jakarta, (1): 30-64 (in Indonesian)

_____, 1983 Marine fishery development in Sumatra. Mar.Fish.Res.Rep.,Jakarta, (28):1-12 (in Indonesian)

Sujastani, T. and S. Nurhakim, 1982 On the potential of lemuru (Sardinella longiceps). In Proceedings of the Seminar on lemuru fishery, Banyuwangi, 18-21 January 1982, pp. 1-12

Sumiono, B., 1983 The length of first maturity stage and sex ratio of Penaeus merguiensis de Man in the Bintuni Bay, Irian Jaya. Mar.Fish.Res.Rep.,Jakarta, (29):41-51 (in Indonesian)

Supongpan, M., 1984 The cephalopod fisheries and resources in the Gulf of Thailand. Paper presented to the Marine Fisheries Seminar, Bangkok, 4-7 September 1984, 20 p. (mimeo) (in Thai)

Supongpan, M. and K. Kongmuag, 1976 Study on the sex ratio of the squids and cuttlefish in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1976):42 p. (in Thai)

_____, 1976a Study on the length-weight relationship of the squids and cuttlefish in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1976):18 p. (in Thai)

Supongpan, S., 1979 Pla-Insi. In Results of the Seminar on Marine Fisheries Development Program. Bangkok, Marine Fisheries Division, Department of Fisheries, Annex 4-7, 10 p. (in Thai)

Supongpan, S. and R. Chayakul, 1979 Population bionomics of the Spanish mackerel, Scomberomorus commerson (Lacépède, 1802) in the Gulf of Thailand. Tech.Rep.Pelagic Fish.Unit Mar. Fish.Div.,Bangkok, (9):31 p.

Supongpan, S., et al., 1982 Light-fishing purse seiners in the Gulf of Thailand, 1981. Tech.Rep. Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (27):57 p. (in Thai)

_____, 1984 A study on age and growth of Stolephorus heterolobus (Rüppel, 1937) in the inner Gulf of Thailand (1970-1973). Tech.Rep.Pelagic Fish.Unit Mar.Fish.Div.,Bangkok, (44):23 p. (in Thai)

Suvapepun, S. and W. Suwanrumpha, 1970 Distribution of copepods in the inner Gulf of Thailand and the western coast of the Gulf of Thailand. Proc.IPFC, 13(2):1-19

Tabtimtai, C., 1968 Relationship between the fecundity and the length and the weight of chub mackerel. Annu.Rep.Pelagic Fish.Unit Mar.Fish.Lab.,Bangkok, (1966-1967):88-101 (in Thai)

Tan, E.O., 1970 Notes on the biology of chub mackerel, Rastrelliger brachysoma (Bleeker), in Manila Bay. In The Kuroshio: a Symposium on the Japan Current, edited by J.C. Marr. Honolulu, East-West Center Press, pp. 479-890

Tan, K.S., 1974 Fisheries biological study of groupers in the South China Sea. In Working papers of trainers, vol.3, 1973. Singapore, SEAFDEC, Marine Fisheries Research Department, pp. 31-61

Tan, S.M. and K.S. Tan, 1974 Biology of the tropical grouper, Epinephelus tauvina (Forskal). 1. A preliminary study on hermaphroditism in E. tauvina. Singapore J.Primary Ind., 2(2): 123-33

Tantiswetratana, C., 1979 Pla-Lung. In Results of the Seminar on Marine Fisheries Development Program organized by Marine Fisheries Division, Department of Fisheries, Bangkok, Annex 4, 5 p. (in Thai)

Taweesith, T., 1979 Anchovies. In Results of the Seminar on Marine Fisheries Development Program organized by Marine Fisheries Division, Department of Fisheries, Bangkok, Annex 4, 10 p. (in Thai)

Termvidchakorn, A., 1983 Studies on the development and distribution of the carangid fishes in the Kuroshio and adjacent regions. Ph.D. Thesis, Ocean Research Institute, University of Tokyo (Unpubl.)

Tham, A.K., 1953 A preliminary study of physical, chemical and biological characteristics of Singapore Straits. Fish.Publ.Colon.Off.Lond., 1(4):65 p.

_____, 1967 A contribution to the study of the growth of members of the genus Stolephorus Lacépède in Singapore Strait. Proc.IPFC, 12(2):1-25

_____, 1967a Study on the growth of fish in Malaysia and Singapore waters. In Proceedings of the Second Symposium on scientific and technological research in Malaysia, pp. 94-104

_____, 1974 Stolephorus resources in the South China Sea. Proc.IPFC, 15(3):182-91

Thubtimsaeng, V., 1978 Observation on the distribution of shrimp larvae in the Gulf of Thailand. Tech. Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (28/1978)

_____, 1978a Distribution of some invertebrate larvae in the Gulf of Thailand. Tech.Rep. Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (27/1978)

Thubtimsang, W., 1977 Observation on the spawning ground and spawning season of economically important shrimps in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1977):48 p. (in Thai)

Thubtimsang, W. and A. Meechukhan, 1976 Observation on the spawning ground and spawning season of economically important shrimp in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1976):41 p. (in Thai)

Thumniyom, D. and T. Charnprasertporn, 1977 Length-weight relationship of monocle bream (Scolopsis taeniopterus) in the area off Cholburi, inner Gulf of Thailand, June 1970 - May 1971. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (2/2520):21 p. (in Thai)

Tiews, K. (ed.), 1976 Fisheries resources and their management in Southeast Asia. Proceedings of the International Seminar on fisheries resources and their management in Southeast Asia, organized by the German Foundation for International Development in cooperation with the Federal Research Board for Fisheries and FAO. Berlin, 19 November to 6 December 1974. Bonn, Druckerei Heinz Lammerich, 522 p.

Tiews, K., S.A. Bravo and I.A. Ronquillo, 1972 On the food and feeding habits of some Philippine shrimps in Manila Bay and San Miguel Bay. Proc.IPFC, 13(3):85-92

Tiews, K., A. Mines and I.A. Ronquillo, 1972 On the biology of Saurida tumbil (Bloch, 1801), Family Synodontidae in Philippine waters. Proc.IPFC, 13(3):100-20

Tiews, K., I.A. Ronquillo and P. Caces-Borja, 1970 On the biology of round scads, Decapterus Bleeker in the Philippine waters. Proc.IPFC, 13(2):82-106

Tiews, K., I.A. Ronquillo and I.M. Sartos, On the biology of anchovies, Stolephorus Lacépède in
1970 the Philippine waters. Proc.IPFC, 13(2):20-48

Tiews, K., et al., On the food and feeding habit of eight species of Leiognathus found in Manila
1972 Bay and San Miguel Bay. Proc.IPFC, 13(3):93-9

Tu, L.V., The fecundity and biological minimum size of five commercial important species in the
1971 South China Sea. Graduation thesis submitted to Marine Fisheries Research Department, SEAFDEC, Singapore, 22 p.

Uchida, R.N., Synopsis of biological data on frigate tuna, Auxis thazard, and bullet tuna,
1981 A. rochei. FAO Fish Synop., (124):63 p. Issued also as NOAA Tech.Rep.NMFS Circ., (436):63 p.

Uktolseja, J.C.B., et al., Summberdaya Perikanan Laut Sulawasi Tengah Tahap. 1. Laporan
1979 Penelitian Perikanan Laut/Mer.Res.Rep.,Jakarta, (2/79) (in Indonesian)

Vadhanakul, P., Spawning season and spawning ground of Nemipterus nematophorus in the Gulf of
1980 Thailand, 1979. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (5/1980):17 p. (in Thai)

_____, Spawning season and spawning ground of Nemipterus mesoprion in the Gulf of Thailand.
1981 Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (2/1981):21 p. (in Thai)

Vadhanakul, S., Preliminary study on the life history of Leiognathus brevirostris and Leiognathus
1976 spp. in the inner Gulf of Thailand. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (2/2519):22 p. (in Thai)

_____, Study on the life history of Lutjanus lineolatus (Rüppel) in the inner Gulf of
1976a Thailand. Tech.Rep.Demersal Fish.Unit Mar.Fish.Div.,Bangkok, (3/2519):17 p. (in Thai)

Vanichkul, P. and V. Hongskul, Length-weight relationships of chub mackerel (Rastrelliger spp.) in
1965 the Gulf of Thailand, 1963. In Reports on mackerel investigations, 1963-1965. Contrib.mar.Fish.Lab.,Bangkok, (4):pp. 162-89 (in Thai)

Vattanachai, S., The identification of fish eggs and larvae obtained from the survey cruise in the
1974 South China Sea. Proc.IPFC, 15(3):111-30

Vibhasiri, A., Penaeid shrimp resources and fisheries in the Gulf of Thailand. Paper presented to
1984 the Marine Fisheries Seminar, 4-7 September 1984. Bangkok, Department of Fisheries, 21 p. (mimeo) (in Thai)

Vibhasiri, A. and K. Kongmuag, Length-weight relationship of three commercially important penaeid
1976 shrimps in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.Unit Mar.Fish.Div.,Bangkok, (1976):13 p. (in Thai)

_____, Sex ratios of penaeid shrimps in the Gulf of Thailand. Annu.Rep.Invertebr.Fish.
1976a Unit Mar.Fish.Div.,Bangkok, (1976):17 p. (in Thai)

Wahyuono, H., et al., Biological parameters of some demersal fish species in the Malacca Strait,
1983 North Sumatra Province. Mar.Fish.Res.Rep.,Jakarta, (26):29-47

Weber, W. and A. Jothy, Observations on the fish Nemipterus spp. (Family: Nemipteridae) in the
1977 coastal waters of East Malaysia. Arch.Fisch.Wiss., (28):109-22

Weng, C.C., Commercial prawns of Peninsular Malaysia. Fish.Bull.Minist.Agric.Malays., (21):43 p.
1978

Wetchagarun, K., Some aspects on the biology of spotfinned bigeye (Priacanthus tayenus) in the
1971 inner part of the Gulf of Thailand (1969-1970). Paper presented to the Second Symposium on marine fisheries, Bangkok, Marine Fisheries Laboratory, 19-20 April 1971, 24 p. (mimeo)

Whitehead, P.J.P., FAO species catalogue. Vol.7. Clupeoid fishes of the world. An annotated and
1985 illustrated catalogue of the herrings, sardines, pilchards, sprats, anchovies and wolf-herrings. Part 1. Chirocentridae, Clupeidae and Pristigasteridae. FAO Fish.Synop., (125) Vo.7, Pt.1:303 p.

Whitehead, P.J.P., FAO species catalogue Vol.7. Clupeoid fishes of the world. Annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, anchovies and wolf-herrings. Part 2. Engraulidae. FAO Fish.Synop., (125) Vol.7, Pt. 2 (in preparation)

Widodo, J., 1976 A check list of the fish collected by Mutiara-4 from November 1974 to November 1975. Contrib.Demersal Fish.Proj.(Indones.-Ger.),Jakarta, (1):47-77

Wongchitsue, B., 1984 Study on the length-weight relationship of round scads (Decapterus spp.) along the west coast of Thailand. Technical Report, Phuket, Phuket Marine Fisheries Station, 15 p. (in Thai)

Wongratana, T., 1968 A checklist of fishes caught during the trawl surveys in the Gulf of Thailand and off the east coast of Malay Peninsula. Contrib.Mar.Fish.Lab.Mar.Fish.Div.,Bangkok, (13):96 p.

Wyrtki, K., 1961 Physical oceanography of the Southeast Asian waters. Naga Rep., (2):195 p.

Yesaki, M., 1982 Biological and environmental observation (tuna). A report prepared for the Pole and Line Tuna Fishing in Southern Thailand Project. Rome, FAO, FI:DP/THA/77/008, Field document 3:50 p.

LIST OF TABLES

		Page
1.	Occurrence of major species groups and species in the region	28
	(1) Coastal pelagic fish	
	(2) Demersal fish	
	(3) Shrimps	
	(4) Cephalopods	
2.	Important biological features and parameters; coastal small pelagic fish in the region	32
	(1) Mackerels	
	(2) Round scads	
	(3) Sardines	
	(4) Anchovies	
	(5) Carangids	
3.	Important biological features and parameters; king mackerels and coastal tunas in the region	41
	(1) King mackerels	
	(2) Coastal tunas	
4.	Important biological features and parameters; demersal fish in the region	43
	(1) Pony fish	
	(2) Red snappers	
	(3) Goat fish	
	(4) Threadfin breams	
	(5) Bigeye snappers	
	(6) Groupers	
	(7) Croakers	
	(8) Lizardfish	
5.	Important biological features and parameters; shrimps in the region	55
	(1) Penaeus spp.	
	(2) Metapenaeus spp.	
	(3) Acetes spp.	
6.	Important biological features and parameters; cephalopods in the region	60
	(1) Squids	
	(2) Cuttlefish	
	(3) Octopuses	

Table 1

Occurrence of major species groups and species in the region
(1) Coastal pelagic fish. See Section 4 for annotation and Sections 2 and 6.1 for references

Family and genus	Number of species involved in genus	Species of commercial value reported from countries									
		Species	Brunei	Burma	Hong Kong	Indonesia	Kampuchea	Malaysia	Philippines	Singapore	Thailand	Viet Nam
FAMILY CARANGIDAE												
Alectis	2											
Alepes	3	A. kalla		+	+						+	
Atropus	1	-										
Atule	1	A. mate		+	+	+	+	+	+		+	+
Carangoides	13	C. malabaricus		+	+			+		+	+	
Caranx	8	-										
Decapterus	8	D. macrosoma				+		+	+		+	
		D. maruadsi			+			+	+		+	
		D. russelli		+		+			+			+
Elagatis	1	-										
Gnathonodon	1	-										
Megalaspis	1	M. cordyla	+	+	+	+	+	+	+	+	+	+
Naucrates	1	-										
Scomberoides	4	S. lysan		+				+			+	
Selar	3	S. boop		+				+				
		S. crumenophthalmus	+	+	+	+	+	+	+	+	+	+
		S. malam		+							+	
Selaroides	1	S. leptolepis	+	+	+	+	+	+	+	+	+	+
Seriola	2	-										
Seriolina	1	S. nigrofasciata		+				+			+	
Trachinotus	3	-										
Trachurus	1	T. japonicus			+							
Ulva	1	-										
FAMILY CLUPEIDAE												
Anodontostoma	2	-										
Dussumieria	1	D. acuta				+		+	+		+	+
Escualosa	1	-										
Herklotsichthys	2	-										
Hilsa	4	-										
Ilisha	5	-										
Nematalosa	3	-										
Opisthopterus	1	-										
Pellona	1	-										
Raconda	1	-										
Sardinella	11	S. fimbriata		+		+		+	+		+	
		S. gibbosa		+		+					+	
		S. leiogaster							+		+	+
		S. longiceps				+						
Ambligaster		A. sirm				+			+		+	
Spratelloides	2	-										
FAMILY ENGRAULIDAE												
Coilea	5	-										
Setipinna	8	-										
Stolephorus	13	S. heterolobus	+	+	+	+	+	+	+	+	+	+
		S. indicus	+	+	+	+	+	+	+	+	+	+
Thrissina	1	-										
Thryssa	8	-										

Table 1 (1) (continued)

Family and genus	Number of species involved in genus	Species of commercial value reported from countries									
		Species	Brunei	Burma	Hong Kong	Indonesia	Kampuchea	Malaysia	Philippines	Singapore	Thailand	Viet Nam
FAMILY SCOMBRIDAE												
Acanthocybium	1	-										
Auxis	2	A. thazard	+	+	+	+	+	+	+	+	+	+
Grammatorcynus	2	-										
Gymnosarda	1	-										
Euthynnus	1	E. affinis										
Rastrelliger	3	R. brachysoma	+	+	+	+	+	+	+	+	+	+
		R. kanagurta	+	+	+	+	+	+	+	+	+	+
Sarda	2	-										
Scomber	2	-										
Scomberomorus	5	S. commerson	+	+	+	+	+	+	+	+	+	+
Thunnus	5	T. tonggol	+	+	+	+	+	+	+	+	+	+

Table 1 (continued)

(2) Demersal fish. See Section 4 for annotation and Sections 2 and 6.2 for references

Family and genus	Number of species involved in genus	Species of commercial value reported from countries									
		Species	Brunei	Burma	Hong Kong	Indonesia	Kampuchea	Malaysia	Philippines	Singapore	Thailand	Viet Nam
FAMILY LEIOGNATIDAE												
Gazza	1	G. minuta		+		+		+	+			
Leiognathus	20	L. bindus		+		+		+	+		+	
		L. blochii							+		+	
		L. brevirostris			+	+			+		+	
		L. equulus		+		+		+	+		+	+
		L. splendens		+		+			+		+	+
Secutor	2											
FAMILY LUTJANIDAE												
Lutjanus	24	L. argentimaculatus		+							+	+
		L. johni		+		+			+		+	
		L. lineolatus				+		+			+	
		L. malabaricus		+	+	+	+	+	+	+	+	
		L. sebae	+			+					+	
		L. sanguineus		+		+						+
		L. vitta				+	+		+		+	
Pristipomoides	4	P. typus									+	
Casio	7	-		+		+			+			
Glabilutjanus	1	-										
FAMILY NEMIPTERIDAE												
Nemipterus	20	N. hexodon				+		+	+		+	
		N. japonicus		+				+	+		+	+
		N. nematophorus				+		+			+	
		N. peronii				+		+			+	
		N. nemurus				+		+				
Scolopsis	11	S. taeniopterus				+		+			+	
Parascolopsis	2	-										

Table 1 (2) (continued)

Family and genus	Number of species involved in genus	Species of commercial value reported from countries									
		Species	Brunei	Burma	Hong Kong	Indonesia	Kampuchea	Malaysia	Philippines	Singapore	Thailand	Viet Nam
FAMILY SCIAENIDAE												
Bahaba	2	-										
Chrysochir	1	-										
Collichthys	2	-										
Dendrophysa	1	D. russelli				+		+	+		+	
Johnieops	3	J. sina				+		+				
		J. dussumieri		+				+			+	
Johnius	6	-										
Kathala	1	-										
Nibea	4	-										
Otolithes	4	O. argenteus		+							+	+
Otolithoides	2	-										
Panna	1	-										
Pennahia	4	P. macrocephalus							+		+	
Protonibea	1	-										
Pterololithes	2	-										
Pseudosciaena	1	-										
FAMILY PRIACANTHIDAE												
Priacanthus	3	P. macracanthus		+	+	+					+	
		P. tayenus	+	+	+	+	+	+	+	+	+	+
FAMILY MULLIDAE												
Upeneus	7	U. bensasi						+			+	+
		U. moluccensis			+				+		+	
		U. sulphureus		+		+			+		+	
Mulloidichthys	1	-										
Parupeneus	6	-										
FAMILY SERRANIDAE												
Cephalopholis	8	-										
Cromiloptes	1	-										
Epinephelus	21	E. awoara			+						+	
		E. megachir						+			+	
		E. sexfasciatus						+	+		+	+
		E. tauvina			+	+			+		+	+
Plectopomus	5	-										
Promicrops	1	-										
Variola	3	-										
Serranus	1	-										
Anyperodon	1	-										
Aethaloperca	1	-										
FAMILY SYNODONTIDAE												
Saurida	9	S. tumbil		+				+	+		+	+
		S. undosquamis				+		+	+		+	

Table 1 (continued)

(3) Shrimps. See Section 4 for annotation and Sections 2 and 6.3 for references

Family and genus	Number of species involved in genus	Species of commercial value reported from countries										
		Species	Brunei	Burma	Hong Kong	Indonesia	Kampuchea	Malaysia	Philippines	Singapore	Thailand	Viet Nam
FAMILY PENAEIDAE												
Penaeus	11	P. indicus		+			+	+	+		+	
		P. latisulcatus				+	+	+	+		+	
		P. merguiensis	+	+	+	+	+	+	+	+	+	+
		P. monodon				+	+	+	+		+	+
		P. semisulcatus			+	+	+	+	+		+	
Metapenaeus	14	M. affinis	+			+		+			+	
		M. brevicornis	+			+	+	+				
		M. ensis	+			+		+			+	
		M. lysianassa	+			+		+				
Parapenaeopsis	13	P. sculptilis	+			+		+				
Trachypenaeus	4	T. fulvus						+	+		+	
Metapenaeopsis	7	M. stridulans				+		+			+	
Atyopenaeus	2	-										
Parapenaeus	1	-										
Solenocera	3	S. subnuda	+			+		+				

Table 1 (continued)

(4) Cephalopods. See Section 4 for annotation and Sections 2 and 6.4 for references

Family and genus	Number of species involved in genus	Species of commercial value reported from countries										
		Species	Brunei	Burma	Hong Kong	Indonesia	Kampuchea	Malaysia	Philippines	Singapore	Thailand	Viet Nam
FAMILY LOLIGINIDAE												
Loligo	4	L. duvauceli			+	+		+	+		+	
		L. edulis			+	+		+	+			
		L. chinensis			+	+					+	+
Sepioteuthis	2	S. lessoniana			+						+	+
Symplectoteuthis	1	-										
Cranchia	1	-										
Doryteuthis	1	-										
Nototodarus	1	-										
FAMILY SEPEIDAE												
Sepia	8	S. pharaonis							+		+	
		S. recurvirostra							+		+	
Sepiella	2	-										
FAMILY OCTOPODIDAE												
Octopus	5[a/]	O. membranaceus									+	
		O. dollfusi									+	
Cistopus	1[b/]	-										

a/ Although three other species are believed to be widely distributed in the region (Chikuni, 1983, Roper, Sweeney and Nauen (1984), much attention has not been paid by fishermen

b/ Cistopus indicus is also believed to be abundantly distributed in the region (Ibid.), but neither academic nor industrial confirmation of this has been made

Table 2

Important biological features and parameters; coastal small pelagic fish in the region
(1) Mackerels. See Sections 5 and 6.1.1 for annotations and broader references

(Body size refers to total length unless specified as FL: fork length or SL: standard length; sexes are combined unless specified as M: male or F: female)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SCOMBRIDAE Rastrelliger brachysoma	Gulf of Thailand[a/]	20–40	15.0	20.95	10–40 mi off Prachuap Suratthani	2–4, 6–8	egg = $9 \times 10^{-8} L^{4.8356}$ 200 000–500 000, 20 000–30 000/batch	10.25	1–3, 7–9	17.5	1:1	k=0.33	Z=1.06	2–3	Phytoplanktons, zooplanktons	$W = 0.006138L^{3.215}$ $M:W = 0.000005732L^{3.1235}$ $F:W = 0.000006578L^{3.1235}$
	Andaman Sea[b/] (Thailand)	–	17.5	–	Koh Yao, Krabi	2–3, 8–9	30 000/batch 97 250–241 832	9.5–12.5	4, 8–10	17.5	1:1.3	–	–	–	Phytoplanktons, zooplanktons, diatoms, copepods	Log W = 1.8874 + 3.214 Log L
	North of Java[c/]	–	–	22.92	–	6–10	–	–	–	17.3	1.3:1	k=0.19	Z=0.88	3–4	–	–
	Malacca Strait[d/] (Malaysia)	–	–	19.6–20.1	–	10–12	20 000–30 000/batch	10.0	1–3	18.5	–	k=0.36–0.44	M=0.38 Z=0.82	–	–	–
	Manila Bay[e/]	–	–	34.0	–	6–2	11 300–119 300	–	–	15.0–16.0	–	k=1.1	M=1184 Z=4.27	–	–	–
	Samar Sea[f/]	–	–	25.0	–	–	–	–	–	–	–	k=1.60	M=2.56 Z=4.49	–	–	–

a/ Boonprakob (1965, 1967, 1972); Tabtimtai (1968); Suchondhamarn et al. (1970); Somjaiwong et al. (1970); Suvapepun and Suwanrumpha (1970)

b/ Boonragsa et al. (1984); Bussarawitch (1984, 1984a, 1984b)

c/ Sujastani (1974)

d/ Pathansali (1961, 1967); Chong and Chua (1974)

e/ Ingles and Pauly (1984); Tan (1970)

f/ Ingles and Pauly (1984)

Table 2 (1) Mackerels (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SCOMBRIDAE Rastrelliger brachysoma (continued)	Andaman Sea[a/] (Burma)	–	19.3FL	23.0FL	–	10–5	20 000–30 000/ batch	13.7FL	2, 5–6	–	1:1.7	–	–	–	–	–
R. kanagurta	Gulf of Thailand[b/]	30–60	16.0	22.9	–	2–4, 7–8	200 000	11.0	5–6	18.6	1:1	–	–	2–3	Phyto-planktons, zoo-planktons, diatoms, copepods	M:W = $0.0000001958L^{3.7653}$ F:W = $0.000009454L^{3.0375}$
	Andaman Sea[c/] (Thailand)	–	—	19.2	–	12.2	25 000/ batch 94 495 263 178	13.0–14.0	5–12	18.67	1:0.93	–	–	–	Phyto-planktons, diatoms, zoo-planktons, crus-taceans, dinofla-gellates	–
	North of Java[d/]	–	11.9–12.4	23.89	–	10–2, 6–9	200 000 500 000	18.8	–	19.0	1:1.1	k=2.78 k=1.63	Z=1.2 Z=0.58	3–4	Fila-mentous algae, zolina, ceratium	M:W = $1.35x10^{-5}L^{2.9927}$ F:W = $2.16x10^{-5}L^{2.9281}$
	Malacca Strait[e/] (Malaysia)	–	16.75	–	–	All around 5–1, 11–4	20 000–30 000–batch	–	–	18.75	–	–	–	–	Phyto-planktons, crus-taceans, copepods, decapods, dinofla-gellates	–

a/ Druzhinin (1968)

b/ Vanichkul and Hongskul (1965); Boonprakob (1967); Tantiswetratana (1979)

c/ Boonragsa et al. (1984); Bussarawitch (1984)

d/ Sujastani (1974); Gafa (1982); Dwiponggo and Pauly (in press)

e/ Pathansali (1961, 1967); Chee (1980)

Table 2 (1) Mackerels (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SCOMBRIDAE R. kanagurta (continued)	Palawan waters[a]	-	-	28.0	-	-	-	-	-	-	-	k=1.55	Z=8.27 M=2.43	-	-	-
	Andaman Sea[b] (Burma)	-	20.47FL	-	-	9-5	-	14.0FL	6-7	-	1:1.39	-	-	-	-	-

a/ Ingles and Pauly (1984)

b/ Druzhinin (1968)

Table 2 (continued)

(2) Round scads. See Sections 5 and 6.1.2 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (montb)							
FAMILY CARANGIDAE Decapterus macrosoma	Manila Bay and Palawan waters[a]	50-90	M:17.7 F:17.6	25.0	Manila Bay to Palawan	11-3	67 000-106 200	7.0-8.0	2-3	-	1:0.99	0.56 cm/month	-	3-6	Crustaceans, zooplanktons, fish, molluscs	$W = 0.005639L^{3.167}$
	Java Sea[b]	-	-	25.4	-	-	-	17.6	-	-	-	k=0.98	Z=6.22	-	-	-
	Gulf of Thailand[c]	30-60	-	-	-	12-5	-	-	-	16.5	1:0.9	-	-	-	-	-

a/ Magnusson (1970); Tiews, Ronquillo and Caces-Borja (1970); Ronquillo (1974)

b/ Dwiponggo and Pauly (in press)

c/ Chullasorn and Yusukswad (1977)

Table 2 (2) Round scads (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY CARANGIDAE *Decapterus maruadsi*	Gulf of Thailand[a/]	30–70	13.2	23.1	Central Gulf	2–3, 7–8	38 000–515 000	5.5–6.5	1–2, 6–8	16.1	1:1.2	k=0.11 1–2 cm/month	–	2–3	crustaceans, copepods	$W = 0.00005L^{2.811}$
	Java Sea[b/]	–	–	26.8	–	–	–	9–14.2	–	18.3	–	k=0.95–1.04	Z=3.5–4.79 M=1.3	–	–	–
	Malacca Strait[c/] (Malaysia)	–	–	–	–	–	–	7.5	7	–	–	–	–	–	–	–
Decapterus russelli	Palawan waters[d/]	36–180	M:16.72 F:16.5	33.0	Palawan to Manila	11–3 1–4	29 000–49 000	8.0–9.0	2–3	16.6	1:1.04	k=0.2 0.6 cm/month	Z=2.62	3–6	Zooplanktons, crustaceans, fish	$W = 0.0098L^{3.0152}$
	Java Sea[e/]	–	–	26.8	–	–	20 000–84 000	14.6	–	–	–	k=1.09	Z=4.09	–	–	–

a/ Chullasorn and Yusukswad (1977); Chantarasri (1980); Cheunpan (1981)

b/ Sadhotomo *et al.* (1983); Atmadja (1982)

c/ Southeast-Asian Fisheries Development Centre (1982)

d/ Tiews, Ronquillo and Caces-Borja (1970); Magnusson (1970); Ronquillo (1974); Ingles and Pauly (1984)

e/ Dwiponggo and Pauly (in press)

Table 2 (continued)

(3) Sardines. See Sections 5 and 6.1.3 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY CLUPEIDAE Sardinella fimbriata	Manila Bay[a]	10–20	10.0–18.0	18.0	–	–	21 000–61 000	4.0	5, 10	–	–	k=0.7	Z=3.38	–	–	–
	Palawan waters[a]	–	–	22.0	–	–	–	–	–	–	–	k=1.15	Z=6.56	–	–	–
	Malacca Strait[b] (Malaysia)	10–35	13.5	17.0	–	–	–	7.0	6	–	–	–	–	–	–	–
Sardinella gibbosa	Gulf of Thailand[c]	15–40	10.0	18.4	Entire coastal zone	All around 3–4, 7–8	–	8.5	4–8, 12–2	13.3	1:2.3	k=0.33	–	1–2	Phyto-planktons	$W = 9.28 \times 10^{-6} L^{3.0047}$
	Java Sea[d]	–	–	19.5	–	–	–	12.9	–	–	–	k=1.2	Z=6.24	–	–	–
Sardinella longiceps	Bali Strait[e]	20–60	12.7	22.25	Bali Strait to Pang Pung Bay	6–7	60 000–70 000	8.9	–	–	–	k=0.865	Z=3.17	4–5	Copepods	–
	Manila Bay[f]	–	–	21.0	–	–	–	8.0	–	–	–	k=1.1	Z=7.37	–	–	–
	Palawan waters[f]	–	–	23.0	–	–	–	–	–	–	–	k=1.1	Z=7.26	–	–	–
Sardinella melanula	Honda Bay (Parawan)[f]	–	–	22.5	–	–	–	–	–	–	–	k=0.70	Z=3.36	–	–	–
Sardinella sirm	Palawan waters[f]	–	–	27.3	–	–	–	10.0	–	–	–	k=0.86	Z=5.36	–	–	–
	Java Sea[d]	–	–	25.8	–	–	–	18.3	–	–	–	k=1.15	Z=4.31	–	–	–

a/ Ronquillo (1960); Ingles and Pauly (1984)

b/ Southeast-Asian Fisheries Development Centre (1982)

c/ Chullasorn (1979a)

d/ Dwiponggo and Pauly (in press)

e/ Dwiponggo and Pauly (in press); Dwiponggo (1974): Dwiponggo and Uktolseja (1974); Ritterbush (1975); Sujastani and Nurhakim (1982)

f/ Ingles and Pauly (1984)

Table 2 (continued)

(4) Anchovies. See Sections 5 and 6.1.4 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY ENGRAULIDAE Stolephorus bataviensis	Manila Bay[a/]	–	M:7.99 F:7.96	10.8	–	All around 2–12	5 000–10 000	–	2–3	6.5–7.0	1:0.0	–	–	–	–	–
	Malacca Strait[b/] (Malaysia) to Strait of Singapore	–	3.0–6.0	9.9	Adjacent offshore area	All around	–	–	1–6	5.0SL	–	0.57 cm/100 days k=0.17	–	–	Zoo-planktons	–
Stolephorus buccaneeri	Manila Bay[a/]	5–30	M:7.31 F:7.29	13.0	–	9–3	7 000–11 000	5.0–6.0	3	6.5	1:08	3.8–4.0 cm/year	–	–	–	–
Stolephorus commersonii	Manila Bay[c/]	5–30	–	11.3	–	2–6	5 000–10 000	–	–	6.0	1:0.5	k=0.96	Z=4.94	–	Zoo-planktons	–
Stolephorus heterolobus	Manila Bay[c/]	–	M:6.73 F:6.78	9.8	–	All around 9–3	–	3.0–4.0	3–4	6.0	1:0.9	0.24 cm/month k=0.95	Z=10.69	–	–	–
	Strait of Singapore[d/]	–	–	–	–	All around	1 000–2 500	1.5SL	–	5.0SL	–	0.57 cm/100 days	–	–	Phyto-planktons, poly-chaetes	–
	Java Sea[e/]	–	–	9.7	–	–	–	2.6	–	–	–	–	–	2–3	Zoo-planktons, phyto-planktons, crus-taceans	
	Gulf of Thailand[f/]	5–45	4.5	8.89	30 mi off Prachuab Nakorn	3–4, 7–9	1 588	2.8–4.0	All around 4–12	6.0	1:1	k=0.198	–	1–1.5	Phyto-planktons	$M:W = 2.064 \times 10^{-6} L^{3.2494}$ $F:W = 7.089 \times 10^{-6} L^{2.9329}$

a/ Tiews, Ronquillo and Santos (1970);

b/ Tham (1967, 1967a, 1974)

c/ Tiews, Ronquillo and Santos (1970); Ingles and Pauly (1984)

d/ Tham (1967a, 1974);

e/ Southeast-Asian Fisheries Development Centre (1983)

f/ Sitthichokpun (1970); Taweesith (1979); Isara (1972); Supongpan et al. (1984)

Table 2 (4) Anchovies (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY ENGRAULIDAE Stolephorus indicus	Malacca Strait[a] (Malaysia)	–	–	–	–	–	–	–	–	11.6–12.5	–	0.195 cm/100 days	–	–	Zooplanktons, fish, squid larvae	–
	Singapore Strait[b]	–	–	19.0SL	–	–	–	–	–	12.0SL	–	k=0.7118 3.0 cm/42 days	–	–	Zooplanktons	–
	Manila Bay[c]	–	M:11.8 F:11.6	14.6	–	All around	9 000–14 000	8.0	1–3, 9	9.0	1:1.04	k=1.08 3.8–4.0 cm/year	Z=4.53		Zooplanktons, phytoplanktons, fish larvae	–
Stolephorus insularis	Singapore Straight[a]	–	–	11.9SL	–	–	–	–	–	5.0SL	–	k=0.17	–	1	–	–
Stolephorus zollingeri	Manila Bay[d]	–	–	10.6	–	–	–	–	–	–	–	k=1.85	Z=12.0	–	–	–

a/ Tham (1967a)

b/ Tham (1953)

c/ Tiews, Ronquillo and Santos (1970); Ingles and Pauly (1984)

d/ Ingles and Pauly (1984)

Table 2 (continued)

(5) Carangids (excluding round scads, Decapterus spp.) See Sections 5 and 6.1.5 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY CARANGIDAE Atule mate	Gulf of Thailand[a]	15–45	16.0	25.8	30 mi off Chumporn Nakorn	3–4	–	5.5–6.5	1–3, 6–9	–	–	0.8 cm/ k=0.107	–	2–3	–	–
	Malacca Strait[b] (Malaysia)	10–50	16.0	28.0	–	–	–	–	–	–	–	–	–	–	–	–
Atropus atropus	Andaman Sea[c] (Burma)	20–60	17.2	20.0	–	1	–	–	–	–	–	–	–	–	Fish	–
Carangoides malabaricus	Andaman Sea[c] (Burma)	20–60	M:19.9 F:19.3	29.0	–	7–9	–	13.1	1	-	1:1.22	–	–	–	Fish crustaceans, squids	–
	Andaman Sea[d] (Thailand)	20–60	14.7	–	–	All around 2–5, 8–9	–	10–11	1, 3, 7	19.5	1:1.08	–	–	–	–	Log W = 4.2246 + 2.8884 Log L
	Malacca Strait[e] (Indonesia)	–	–	–	–	–	–	–	–	–	–	–	–	–	–	M:W = $0.2727L^{2.9844}$ F:W = $0.0125L^{2.9135}$
	Malacca Strait[f] (Malaysia)	18–45	15.2	–	–	–	–	–	–	–	–	–	–	–	–	–
	South China Sea[g] (Singapore)	–	–	–	–	–	–	–	–	13.0	1:1.4	–	–	–	Fish	–
Elagatis bipinnulatus	Moro Gulf[h]	–	–	97.5	–	–	–	–	7	13.0	1:1.4	k=0.6	Z=1.19 M=0.92	4–5	–	–

a/ Piyathirathitivorakul (1983)

b/ Aglen et al. (1981)

c/ Druzhinin and Hliang (1972)

d/ Bhatia et al. (1979)

e/ Wahyuono et al. (1983)

f/ Latiff et al. (1974)

g/ Tu (1971)

h/ Ingles and Pauly (1984)

Table 2 (5) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
Megalaspis cordyla	Gulf of Thailand[a/]	20–50	22.0	28.8	–	12–5, 8–11	–	11.5–10.5	5, 9	–	1:0.8	1.2 cm/month k=0.2	–	–	Fish, crustaceans	$W = 0.144L^{2.9785}$
	Andaman Sea[b/] (Thailand)	–	–	34.2	–	3–4	–	15.2	9–12	28.0	1:1.24	–	–	–	–	–
Selar crumenophthalmus	Manila Bay[c/]	–	–	36.5	–	–	–	–	–	–	–	k=0.89	Z=2.91 M=1.57	–	–	–
	Java Sea[d/]	–	–	29.5	–	–	–	17.8	–	–	–	k=1.25	Z=5.56	–	–	–
	South China Sea[e/] (Singapore)	–	–	–	–	–	82 000–141 000	–	–	–	1:0.76	–	–	–	–	–
	Gulf of Thailand[f/]	30–60	20–25	–	–	–	–	10.0	–	19.4	1:1.3	–	–	–	–	–
Selaroides leptolepis	Manila Bay[c/]	–	–	29.0	–	–	–	–	–	–	–	k=0.80	Z=2.76	–	–	–
	Visayan Sea[g/]	20–140	3.5–18.5	19.5–23.0	–	8–9	–	3.5–6.0	1–3, 7, 10	M:15.2 F:15.4	1:0.93	k=1.15	Z=8.64	–	–	$W = 0.0063L^{3.19}$
	Gulf of Thailand[h/]	20–50	12.0	19.2 21.0	–	All around 3, 7–8	–	4.0–5.5	6, 11	F:15.4	1:1.02	k=0.128	–	–	Zooplanktons, phytoplanktons, molluscs	M:Log W = 3.257 Log $L^{-5.567}$ F:Log W = 3.629 Log $L^{-6.369}$
	South China Sea[i/] (Indonesia	30–40	–	18.6	–	–	–	10.8	–	–	1:1	k=1.1	–	–		$M:W = 0.0438L^{3.3065}$ $F:W = 0.0104L^{3.0429}$
	Java Sea[d/]	–	–	25.4	–	–	–	17.6	–	–	–	k=0.8	Z=6.22	–	–	–

a/ Nupech et al. (1983); Supongpan et al. (1982)

b/ Bhatia et al. (1979)

c/ Ingles and Pauly (1984)

d/ Dwiponggo and Pauly (in press)

e/ Tu (1971)

f/ Department of Fisheries, Thailand (1984)

g/ Ingles and Pauly (1984); Arce (1981)

h/ Naiyanetr (1963); Nupetch (1984)

i/ Wahyuono et al. (1983); Sudradjat and Nugroho (1983)

Table 3

Important biological features and parameters; king mackerels and coastal tunas in the region
(1) King mackerels. See Sections 5 and 6.1.6 for annotations and broader references

(Body size refers to total length unless specified as FL: fork length or SL: standard length; sexes are combined unless specified as M: male or F: female)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (montb)		Size (cm)	Season (month)							
FAMILY SCOMBRIDAE *Scomberomorus commerson*	Gulf of Thailand[a/]	20–60	50.0	92.0	–	2–3, 6–9	500 000–3 800 000	11.0–21.0	3–5, 7–10	58.6	1:1.6	k=0.12 3.4 cm/months	–	4–5	Fish, molluscs, crustaceans	$W = 0.01302L^{2.8843}$
	Visayan Sea[b/]	–	–	49.0	–	–	–	–	–	–	–	k=0.7	Z=1.49 M=1.23	–	–	–

a/ Chullasorn, Chotiyaputta and Chayakul (1973); Supongpan and Chayakul (1979)

b/ Ingles and Pauly (1984)

Table 3 (continued)

(2) Coastal tunas. See Sections 5 and 6.1.7 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SCOMBRIDAE Auxis thazard	Indonesia[a]	–	–	51.5	–	–	–	23.6	–	–	–	k=1.0	Z=4.03	–	–	–
	Bohol Sea[b]	–	–	47.0	–	–	–	–	–	–	–	k=0.73	Z=2.84	–	–	–
	Moro Gulf[b]	–	–	63.5	–	–	–	–	–	–	–	k=0.72	M=1.28 Z=4.52	–	–	–
	Gulf of Thailand[c]	20 ~	35.0	–	–	4–6, 8–9	–	19.0 27.0	8–11 2, 4–5	34.1	1:1	–	–	3–4	–	$W = 0.00002L^{2.99}$
	Andaman Sea (Thailand)[d]	20 ~	33.5	44	–	2–3, 10–12	–	–	–	38.0	1:1.25	1.4 cm/month k=0.45	–	3–4	–	$W = 0.000011L^{3.119}$
Euthynnus affinis	Gulf of Thailand[c]	20 ~	37.0	–	–	1–3, 6–7	1 730 000	21.0 26.0	2–4, 6.12	37.5	1:1	–	–	–	–	$W = 0.000015L^{2.979}$
	Andaman Sea (Thailand)[d]	20 ~	37.2	64	–	2–4, 9–11	–	–	–	46.0	1:1.3	k=0.63 1.2 cm/month	4	–	–	$W = 0.000015L^{3.034}$
	Philippines[e]	–	42	80	–	All around 4–7	–	–	–	–	–	–	–	–	–	–
Thunnus tonggol	Gulf of Thailand[c]	20 ~	38.5	–	–	3–5, 7–12	1 400 000	22.0–26.0	1–2, 4–6	39.6	1:1	1.5 cm/month	–	4	–	$W = 0.000021L^{2.979}$
	Andaman Sea (Thailand)[d]	–	43.5	56	–	2–4, 8–9	–	–	9–12	44.0	–	1.4 cm/month	–	4	–	$W = 0.000022L^{2.947}$

a/ Dwiponggo and Pauly (in press)

b/ Ingless and Pauly (1984)

c/ Klinmuang (1978, 1981); Cheunpan (1984)

d/ Yesaki (1982)

e/ Ronquillo (1963)

Table 4

Important biological features and parameters; demersal fish in the region
(1) Pony fish. See Sections 5 and 6.2.1 for annotations and broader references

(Body size refers to total length unless specified as FL: fork length or SL: standard length; sexes are combined unless specified as M: male or F: female)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY LEIOGNATHIDAE *Gazza minuta*	Honda Bay, Palawan[a/]	–	–	17.5	–	–	–	5.5	–	–	–	k=0.97	Z=6.62 M=2.03	–	–	–
	San Miguel Bay[a/]	–	–	14.0	–	–	–	–	–	–	–	k=1.1	Z=8.85 M=2.23	–	–	–
Leiognathus bindus	Manila Bay[b/]	–	–	9.3	–	–	–	–	–	–	–	k=1.25	Z=5.35 M=2.88	–	Zooplanktons, fish eggs and larvae, crustaceans	–
	Java Sea[c/]	–	–	12.5	–	÷	–	5.21	–	–	–	k=1.37	Z=8.36	–	–	–
	Gulf of Thailand[d/]	10–50	5.1	11.0	–	5, 8–10, 12	–	3.0, 3.5, 5.0	1, 8, 7	–	–	0.4–0.5 cm/month	–	–	–	–
Leiognathus blochii	Manila Bay[b/]	–	–	12.8	–	–	–	–	–	–	–	k=1.21	Z=4.87 M=2.58	–	Zooplanktons, fish eggs and larvae, crustaceans	–
Leiognathus brevirostris	Gulf of Thailand[e/]	10–50	M:10.0 F:10.5	M:12.0 F:13.5	–	All around 7–9, 11–2, 5	–	M:9.1 F:9.7	–	–	1:1	–	–	–	Zooplanktons, fish eggs and larvae, crustaceans	–

a/ Ingles and Pauly (1984)

b/ Ingles and Pauly (1984); Tiews *et al.* (1972)

c/ Dwiponggo and Pauly (in press)

d/ Amaragsa (1968)

e/ Vadhanakul (1976)

Table 4 (1) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY LEIOGNATHIDAE Leiognathus brevirostris (continued)	Java Sea[a/]	–	–	12.0	–	–	–	7.1	–	–	–	k=0.95	Z=2.79	–	–	–
	Manila Bay[b/]	10–60	M:10.06 F:10.55	M:12.0 F:13.5	–	7–2, 10	–	M:8.0 F:8.5	2, 9	–	1:0.9	k=1.69	Z=12.0 M=3.07	–	Zooplanktons, fish eggs and larvae, crustaceans, molluscs	–
Leiognathus daura	Manila Bay[b/]	–	–	9.5	–	–	–	–	–	–	–	k=1.54	Z=8.1 M=3.24	–	Fish eggs, diatoms, copepods	
Leiognathus equulus	Andaman Sea (Burma)[c/]	20–60	M:15.3 F:16.6	22.0	–	6–9	–	11.1	9–1	–	1:0.6	–	–	–	–	–
	Java Sea[a/]	–	–	21.5	–	–	–	13.4	–	–	–	k=1.5	Z=5.68	–	–	–
	Samar Sea[b/]	–	–	24.8	–	–	–	–	–	–	–	k=1.28	Z=5.97 M=2.22	–	Benthic animals, fish eggs	–
Leiognathus hastatus	Manila Bay[b/]	–	15.5	22.0	–	1–9	–	–	–	–	1:0.6	–	–	–	Zooplanktons, fish eggs and larvae, crustaceans, molluscs	
Leiognathus leuciscus	Java Sea[a/]	–	–	13.5	–	–	–	4.76	–	–	–	k=1.8	Z=6.15	–	–	–
	Gulf of Thailand[d/]	–	7.0	12.0	–	1, 4, 6, 10	–	4.5–6.0	2, 3 8, 9	–	–	0.5–0.6 cm/month	–	–	–	–
	Manila Bay[b/]	–	7.0	12.0	–	1, 4 7, 10	–	4.5	2, 3 8, 9	–	–	k=1.3	Z=7.29	–	Diatoms, fish eggs, benthic animals	

a/ Dwiponggo and Pauly (in press)

b/ Ingles and Pauly (1984); Tiews et al. (1972)

c/ Druzhinin and Hliang (1972)

d/ Amaragsa (1968)

Table 4 (1) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY LEIOGNATHIDAE Leiognathus lineolatus	Malacca Strait[a/] (Indonesia)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	M:W = $0.0263L^{2.754}$ F:W = $0.0668L^{2.412}$
	Gulf of Thailand[b/]	10-50	-	-	-	10-6(6)	-	8.5, 7, 8	2, 4, 6	-	-	-	-	-	-	-
	Manila Bay[c/]	-	-	11.1	-	-	-	-	-	-	-	k=1.22	Z=7.4 M=2.69	-	Diatoms, crustaceams	-
Leiognathus splendens	Java Sea[d/]	-	-	16.9	-	-	-	6.23	-	-	-	k=1.1	Z=4.0	-	-	-
	Samar Sea[c/]	-	-	15.0	-	-	-	-	-	-	-	k=0.72	Z=3.77	-	Crustaceans, diatoms, fish eggs	-
	Manila Bay[e/]	-	-	12.6	-	-	-	-	-	-	-	k=0.74	Z=5.28 M=1.86	-	-	-
Leiognathus ruconius	Manila Bay[c/]	-	M:16.6 F:17.2	M:21.0 F:21.0	-	All around (2, 8)	-	-	-	-	1:0.81	k=1.43 (1-2)	Z=9.54 M=3.21	-	Crustaceans, fish, molluscs, polychaeta	Log W = 2.755 Log L-4.312
Secutor insidiater	Manila Bay[c/]	-	-	10.1	-	-	-	-	-	-	-	k=1.42	Z=7.02 M=3.04	-	Zooplanktons, diatoms, fish eggs, crustaceans	-

a/ Wahyuono et al. (1983)

b/ Vadhanakul (1976)

c/ Ingles and Pauly (1984); Tiews et al. (1972)

d/ Dwiponggo and Pauly (in press)

e/ Ingles and Pauly (1984)

Table 4 (continued)

(2) Red snappers. See Sections 5 and 6.2.2 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY LUTJANIDAE *Lutjanus johni*	South of Kalimantan[a] (Indonesia)	30–60	–	78.2	–	–	–	–	–	–	1.05:1	k=0.21	M=0.61	–	–	$W = 1.7353 \times 10^{-5} L^{2.8885}$
	Andaman Sea[b] (Burma)	40–60	M:52.0 F:54.0	–	–	2–9(9)	–	–	–	–	1:0.7	–	–	–	Fish, crustaceans	
Lutjanus lineolatus	Gulf of Thailand[c]	20–50	13.9	M:18.0 F:19.0	–	10–6	–	8.0	1–4	–	1:1	1–2 cm/month	–	–	Fish, shrimps, squids	$M:W = 0.0263L^{2.754}$ $F:W = 0.0668L^{2.412}$
	Malacca Strait[d] (Indonesia)	–	–	–	–	–	–	–	–	–	1:0.72	–	–	–	–	$M:W = 0.0263L^{2.754}$ $F:W = 0.0668L^{2.412}$
	East coast of Peninsular Malaysia[e]	20–60	15.0	22.0	–	–	–	4.5	6	–	–	–	–	–	–	–
Lutjanus malabaricus	Andaman Sea[f] (Burma)	40–60	M:38.5 F:38.7	60.5	–	7–9	–	21.0	9	–	1:1.2	–	–	–	Fish, molluscs, crustaceans	
Lutjanus sanguinius	South China Sea[g] (Singapore)	–	18.0–66.0	–	–	–	374 000–2 433 000	–	–	32.0	1:0.96	–	–	–	Fish, molluscs, crustaceans	$W = 0.066L^{2.765}$
	South China Sea[h] (Hong Kong)	40–100	–	–	–	3–4	–	–	–	–	–	–	–	–	–	–

a/ Gafa and Sadhotomo (1983)

b/ Druzhinin and Hliang (1972)

c/ Vadhanakul (1976a); Intong (1980)

d/ Wahyuono *et al.* (1983)

e/ Aglen *et al.* (1981)

f/ Druzhinin and Hliang (1972)

g/ Kangwankij (1970); Tu (1971)

h/ Lai-shing (1968)

Table 4 (continued)

(3) Goat fish. See Sections 5 and 6.2.3 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY MULLIDAE Upeneus moluccensis	Samar Sea[a/]	–	–	22.5	–	–	–	–	–	–	–	k=1.1	Z=4.02 M=2.06	–	–	–
	South China Sea[b/] (Hong Kong)	50–110	–	17.5	–	3–9	–	–	–	–	–	k=0.9	–	4	Crustaceans, molluscs, polychaeta	–
Upeneus sulphureus	Java Sea[c/]	30–70	–	25.0	–	–	–	–	–	–	–	k=2.24	–	–	–	$W = 1.6154 \times 10^{-5} L^{2.9768}$
	Andaman Sea[d/] (Burma)	31–60	M:14.6 F:13.0	18.0	–	3–9	–	10.0–11.0	2, 9	–	1:0.7	–	–	–	Crustaceans, fish	–
	Samar Sea[e/]	–	–	19.5	–	–	–	–	–	–	–	k=1.2	Z=6.96 M=2.27	–	–	–
	San Miguel Bay[e/]	–	–	15.3	–	–	–	–	–	–	–	k=1.05	M=2.23	–	–	–
Upeneus vitatus	Manila Bay[e/]	30–60	–	24.5	–	–	–	–	–	–	–	k=0.71	Z=5.18 M=1.51	–	–	–

a/ Lai-shing (1968)

b/ Lee (1974)

c/ Badrudin (1978a)

d/ Druzhinin and Hliang (1972)

e/ Ingles and Pauly (1984)

Table 4 (continued)

(4) Threadfin bream. See Sections 5 and 6.2.4 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month		Size (cm)	Season (month)							
FAMILY NEMIPTERIDAE Nemipterus hexodon	Gulf of Thailand[a/]	10–40	M:16.46 F:15.12	M:27.3 F:24.4	–	All around 1–4, 6–8	–	11.0, 12.0	5, 9, 11	–	1:0.97	(1–2)	–	–	Fish, crustaceans	M:W=0.1161L$^{3.04}$ F:W=0.0176L$^{2.924}$
	Java Sea[b/]	–	–	24.5	–	–	–	10.4	–	–	–	k=0.6	Z=2.54	–	–	–
Nemipterus japonicus	Gulf of Thailand[c/]	20–50	–	M:25.6 F:23.3	–	–	–	10.0	–	–	–	M:k=0.1599 F:k=0.1207	M:Z=5.482 F:Z=4.814	–	–	–
	South China Sea[d/] (Hong Kong)	27–46	–	–	–	3–5	–	–	–	–	–	–	–	–	–	–
	Andaman Sea[e/] (Burma)	31–40	M:20.3 F:17.9	27.0	–	–	–	14.0	1	–	1:0.37	–	–	–	–	–
	Java Sea[b/]	–	–	23.5	–	–	–	10.5	–	–	–	k=0.7	Z=2.24	–	–	–
	Manila Bay[f/]	–	–	30.0	–	–	–	–	–	–	–	k=0.73	Z=3.31 M=1.41	–	–	–
Nemipterus nematophorus	Gulf of Thailand[g/]	15–50 (>40)	M:13.34 F:11.67	M:25.84 F:17.8	–	1–4, 8, 11	–	–	–	11.7	1:0.85	M:k=0.1436 F:k=0.2275	–	–	–	–
	Lingayen Gulf[f/]	–	–	22.0	–	–	–	–	–	–	–	k=0.43	Z=1.48 M=1.12	–	–	–
	Visayan Sea[f/]	–	–	27.0	–	–	–	–	–	–	–	k=0.65	Z=3.38 M=1.39	–	–	–

a/ Rugvichai (1977)

b/ Dwiponggo and Pauly (in press)

c/ Amornchairojkul and Boonwanich (1982)

d/ Lai-shing (1968)

e/ Druzhinin and Hliang (1972)

f/ Ingles and Pauly (1984)

g/ Vadhanakul (1980); Boonwanich (1981)

Table 4 (4) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY NEMIPTERIDAE Nemipterus virgatus	South China Sea[a/]	69–128	–	–	–	3–5	–	–	–	–	–	–	–	–	–	–
Nemipterus mesoprion	Gulf of Thailand[b/]	30–60	M:13.0 F:11.1	M:19.51 F:15.52	–	2–4	–	6.5–7.0	3, 5, 6	–	1:1.1	M:k=0.179 (1.08) F:k=0.224 (0.85)	–	–	Fish, molluscs, crustaceans	M:W = $0.18\times10^{-5}L^{2.93}$ F:W = $7.8\times10^{-5}L^{3.10}$
	Java Sea[c/]	–	–	21.5	–	–	–	10.8	–	–	–	k=0.8	Z=3.25	–	–	–
Nemipterus peronii	Gulf of Thailand[d/]	30–60	M:22.9 F:21.6	M:27.5 F:27.0	–	2–4	–	15.2, 15.5, 15.7	3, 7, 9, 12	–	1:0.84	–	–	–	Worms, fish, squids, crustaceans	M:W = $0.0122L^{2.988}$ F:W = $0.0199L^{3.004}$
Nemipterus ovenii	Visayan Sea[e/]	20–140	14.0	25.0	–	–	–	5.5–8.0	2, 10, 12	–	1:1.4	k=0.42	Z=1.52 M=1.06	–	–	$W=0.00895L^{3.26}$
Nemipterus delagoae	Malacca Strait[f/] (Indonesia)	–	–	–	–	–	–	–	–	–	1:1.24	–	–	–	–	$W=0.0201L^{2.7735}$
Scolopsis taeniopterus	Gulf of Thailand[g/]	10–40	M:21.7 F:14.6	M:27.0 F:25.0	–	All around 12–1, 4–8	–	–	6–7, 11–12	–	1:0.8	–	–	–	Crustaceans, fish	M:W = $1.08\times10^{-4}L^{2.6201}$ F:W = $6.17\times10^{-5}L^{2.718}$

a/ Lai-shing (1968)

b/ Boonwanich (1980); Vadhanakul (1981)

c/ Dwiponggo and Pauly (in press)

d/ Isarankura (1968); Isarankura and Pariyanond (1963)

e/ Ingles and Pauly (1984); Arce (1981)

f/ Wahyuono et al. (1983)

g/ Thumniyom and Charnprasertporn (1977)

Table 4 (continued)

(5) Bigeye snappers. See Sections 5 and 6.2.5 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY PRIACANTHIDAE Priacanthus macracanthus	North of Java[a]	–	7.5–22.7	26.0	–	2–4, 9–11	–	12.7	–	–	–	k=1.36	Z=5.42 M=3.45	4–5	–	$W=8.7\times10^{-5}L^{2.667}$
Priacanthus tayenus	Gulf of Thailand[b]	40–50	M:27.0 F:25.0	–	–	All around 1–3	56 000–152 000	11.0, 12.0, 10.5	3, 5, 10, 12	14.0	1:1	2.0 cm/month	–	–	Crustaceans, fish, squid	$M:W=3.16\times10^{-6}L^{2.919}$ $F:W=2.606\times10^{-6}L^{2.891}$
	Andaman Sea[c] (Thailand)	–	–	11.9	–	10–4	–	–	–	15.0	1:0.72	–	–	–	–	$\text{Log } W = -3.3815 + 2.4607 \text{ Log} L$
	Samar Sea[d]	–	–	29.0	–	–	–	–	–	–	–	k=1.25	Z=8.95 M=8.09	–	–	–

a/ Nugroho et al. (1983)

b/ Chomjurai (1970); Wetchagarun (1971)

c/ Chantawong et al. (1984)

d/ Ingles and Pauly (1984)

Table 4 (continued)

(6) Groupers. See Sections 5 and 6.2.6 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month		Size (cm)	Season (month)							
FAMILY SERRANIDAE *Epinephelus sexfasciatus*	South China Sea[a/] (Singapore)	30–60	–	86.5	–	1–4, 8–9	2.0–2.7 million	–	–	45–50	–	k=0.14	–	–	–	–
	Gulf of Thailand[b/]	20–70	–	–	–	–	–	–	–	–	–	–	–	–	Fish, crustaceans, molluscs, polychaetae	–

a/ Tan (1974)

b/ Intong (1980a)

Table 4 (continued)

(7) Croakers. See Sections 5 and 6.2.7 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at forst maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SCIAENIDAE Dendrophysa russelli	San Miguel Bay[a/]	–	–	17.5	–	–	–	–	–	–	–	k=0.95	Z=5.32 M=2.01	–	–	–
	Gulf of Thailand[b/]	–	M:11.97 F:11.52	M:16.0 F:15.5	–	All around 12-2	–	–	3, 6, 9, 11, 1	–	1:1	1 cm	–	–	–	$W = 1.259 \times 10^{-7} L^{2.9927}$
Johnius dussumieri	Andaman Sea[c/] (Burma)	20–60	M:21.8 F:21.2	25.0	–	3–9	–	–	–	–	–	–	–	–	–	–
Otolithes lateoides	Andaman Sea[c/] (Burma)	20–60	M:20.3 F:19.7	25.5	–	1–9	–	–	–	–	1:0.6	–	–	–	Fish, crustaceans	–
Otolithes ruber	San Miguel Bay[a/]	–	–	29.5	–	–	–	–	–	–	–	k=0.44	Z=1.71 M=1.025	–	–	–
Pennahia macrophthalmus	Manila Bay[a/]	–	–	26.5	–	–	–	–	–	–	–	k=1.4	Z=5.55 M=2.30	–	–	–
	San Miguel Bay[a/]	–	–	20.0	–	–	–	–	–	–	–	k=0.6	Z=2.28 M=1.43	–	–	–

a/ Ingles and Pauly (1984)

b/ Phettongkam and Thasananukulkij (1972)

c/ Druzhinin and Hliang (1972)

Table 4 (continued)

(8) Lizardfish. See Sections 5 and 6.2.8 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SYNODONTIDAE Saurida elongata	Gulf of Thailand[a/]	10–20	M:26.3 F:30.6	M:37.7 F:41.6	< 30 mi	1–3, 8–9	–	–	5–7, 11	–	–	M:K=0.103 1.4 cm/month F:K=0.099 1.5 cm/month	M:Z= 5.622 F:Z= 5.278	–	–	M:W = $5.644 \times 10^{-6} L^{3.054}$ F:W = $6.565 \times 10^{-6} L^{3.024}$
Saurida tumbil	Gulf of Thailand[b/]	–	M:26.31 F:30.56	M:34.0 F:40.0	–	1–3	–	–	5–7, 12	–	1:1	–	–	–	–	–
	Andaman Sea[c/] (Burma)	21–60	29.5	38.0	–	2.8	–	–	–	–	1:1.7	–	–	–	–	–
	Manila Bay[d/]	–	37.0	All year round	–	–	28 000	7.0–8.0	11–2	M:17.0–19.0 F:19.0–21.0	1:0.8	K=1.03 0.66 cm/month	Z=4.83 M=1.71	–	Fish	–
	Visayan Sea[d/]	–	41.0	2–5, 7–9	–	–	–	–	–	–	–	k=0.70	Z=2.22 M=1.30	–	–	–
Saurida undosquamis	Gulf of Thailand[e/]	41–50	M:17.8 F:18.31	M:36.5 F:26.5	–	12–1, 5–9	–	12.0–14.0	6, 12, 2	–	1:0.57	2.0 cm/month	–	–		$W = 0.000002929 L^{3.163}$
	Java Sea and Malacca Strait[f/] (Indonesia)	–	–	33.5	–	–	–	15.15	–	–	1:31	k=0.95	Z=2.9	–	–	M:W = $0.0002 L^{4.0843}$ F:W = $0.0002 L^{4.0759}$
	Visayan Sea[g/]	–	–	30.5	–	–	–	–	–	–	–	k=0.80	Z=4.07	–	–	–

a/ Charnprasertport (1979)

b/ Chomdej (1966)

c/ Druzhinin and Hliang (1972)

d/ Tiews, Mines and Ronquillo (1972); Ingles and Pauly (1984)

e/ Intong (1971, 1976); Shinoda and Intong (1977)

f/ Wahyuono et al. (1983); Dwiponggo and Pauly (in press)

g/ Ingles and Pauly (1984)

Table 4 (8) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SYNODONTIDAE Saurida longimanus	Java Sea[a/]	-	-	20.5	-	-	-	11.28	-	-	-	k=1.0	Z=2.19	-	-	-
Saurida micropectolaris	Java Sea and Malacca Strait[b/] (Indonesia)	-	-	42.0	-	-	-	13.84	-	-	1:0.56	k=0.875	Z=3.73 M=1.5	-	-	M:W = $0.0028L^{3.2812}$ F:W = $0.0189L^{2.7424}$

a/ Dwiponggo and Pauly (in press)

b/ Dwiponggo and Pauly (in press); Wahyuono et al. (1983)

Table 5

Important biological features and parameters; shrimps in the region
(1) Penaeus spp. See Sections 5 and 6.3 for annotations and broader references

(Body size refers to total length unless specified as CL: carapace length or g: body weight in grammes; sexes are combined unless specified as M: male or F: female)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY PENAEIDAE Penaeus indicus	Malacca Strait[a]	–	–	–	–	All around 1–4, 7–10	124 000 1 500 000	–	–	–	–	–	–	–	Phytoplanktons, polychaeta, feraminifera	–
	South China Sea[b] (Brunei)	–	–	20.0 CL (60 g)	–	4, 9–11	–	–	11, 4	3.0 CL	–	3.2 cm CL/ 6 months	–	–	–	Log W = 2.922 Log C–0.16
Penaeus merguiensis	Gulf of Thailand[c]	15–30	–	–	–	1–3, 9–12	129 650–960 950	–	–	13.0–14.2	–	–	–	–	Polycheata, fish, copepods, euphausia	M:W = $0.0000100L^{2.963}$ F:W = $0.00000497L^{3.113}$
	Andaman Sea[d] (Thailand)	–	–	–	–	7–8	–	–	–	13.0	–	–	–	–	–	–
	South China Sea[b] (Brunei)	–	4.85 CL	–	Inshore	All around 6, 9	–	2.2 CL	10, 2	–	1:1	2–3 g/ month, 1.7 cm/ 97 days	–	–	–	M:Log W = 3.04 Log C–0.033 F:Log W = 3.06 Log C–0.04
	Manila Bay[e]	–	–	–	–	–	–	–	–	–	–	–	–	–	Zoo/phytoplanktons, bentic feraminifera, ostracods, copepods	–
	San Miguel Bay[e]	–	–	–	–	–	–	–	–	–	–	–	–	–	Same as above	–

a/ Pathansali et al. (1974)

b/ Currie (1982)

c/ Hutasing (1974); Vibhasiri and Kongmuag (1976, 1976a); Thubtimsang (1977); Thubtimsang and Meechukhan (1976); Thubtimsaeng (1978, 1978a)

d/ Kungvankij et al. (1973, 1973a)

e/ Tiews, Bravo and Ronquillo (1972)

Table 5 (1) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY PENAEIDAE Penaeus merguiensis (continued)	Java Sea[a/]	-	6-12	24.6	-	3, 4, 8, 11, 12	-	-	-	2.6 CL 12.5	1:1.18	k=1.2	Z=5.98 M=3.16	-	-	-
	South China Sea[b/] (Sabah, Malaysia)	-	-	-	-	-	-	-	-	3.38 CL	1:1.18	-	-	-	-	-
Penaeus canaliculatus	Malacca Strait[b/] (Malaysia)	-	-	-	-	-	200 000	-	-	-	-	-	-	-	-	-
	Manila Bay[c/]	> 40	-	-	-	-	-	-	-	-	-	-	-	-	Zoo/phytoplanktons, benthic feraminifera, ostracods, copepods, fish eggs	-
	San Miguel Bay[c/]	-	-	-	-	-	-	-	-	-	-	-	-	-	Same as above	-
Penaeus japonicus	Gulf of Thailand[d/]	30-39	17.7	25.0	-	1-3, 10-12	730 000	11.0	-	15.0-15.6	0.74:1	-	-	-	Phyto/zooplanktons, molluscs, cephalopods	M:W = $0.000007L^{3.0087}$ F:W = $0.000004L^{3.1213}$
Penaeus latisulcatus	Gulf of Thailand[e/]	10-19	12.9	20.9	-	All around 1-3, 7-8	257 889-1 009 459	7.0	-	14.0	1:1	-	-	-	Shrimps, crabs, cephalopods, molluscs	M:W = $0.0000712L^{2.5703}$ F:W = $0.0000149L^{2.9018}$
Penaeus longistylus	Gulf of Thailand[f/]	10-19	13.9	22.4	-	-	-	-	-	-	-	-	-	-	-	-
Penaeus monodon	Gulf of Thailand[g/]	30-39	24.5	31.9	-	All around 1-3, 6-7	100 000-300 000	-	-	20.0	1:1	-	-	-	Polychaeta, decapeda, euphausiacea	

a/ Martosubroto (1977); Sumiono (1983); Adisusilo (1983)

b/ Pathansali et al. (1974)

c/ Tiews, Bravo and Ronquillo (1972)

d/ Thubtimsang and Meechukhan (1976); Chaitiamvong (1980)

e/ Thubtimsang and Meechukhan (1976); Chaitiamvong (1980); Chaitiamvong and Thaiprayoon (1979)

f/ Vibhasiri (1984)

g/ Thubtimsang (1977); Thubtimsaeng (1978, 1978a)

Table 5 (1) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY PENAEIDAE Penaeus monodon (continued)	Andaman Sea[a] (Thailand)	-	-	-	-	2, 8	-	-	-	20.0	-	-	-	-	-	-
	Malacca Strait[b] (Malaysia)	-	-	-	-	-	300 000	-	-	-	-	-	-	-	-	-
	Arafura Sea[c] (Indonesia)	-	-	43.0	-	-	-	-	-	21.0	-	k=1.63	Z=5.05	-	-	-
	South China Sea[d] (Brunei)	16	M:8-64 g F:11-175 g	-	-	-	-	-	-	-	-	-	-	-	-	-
Penaeus semisulcatus	Gulf of Thailand[e]	35-40	147	-	-	5-6	178 438-855 579	-	-	15.1	0.94:1	-	-	-	Crustaceans, molluscs, squids	M:W = $0.00000945L^{2.9947}$ F:W = $0.00000216L^{3.2931}$
	Andaman Sea[a] (Thailand)	-	-	-	-	2, 8	-	-	-	16	-	-	-	-	-	-
	Malacca Strait[b] (Malaysia)	-	-	-	-	-	-	-	-	-	-	-	-	-	Zoo-phyto-planktons, benthos, feraminifera	
	Manila Bay[f]	-	-	-	-	-	-	-	-	-	1:0.5	-	-	-	Zoo/phyto-planktons, benthos, feraminifera, molluscs, echinoderma-ta, fish eggs	-

a/ Kungvankij et al. (1973a)

b/ Pathansali et al. (1974)

c/ Dwiponggo and Pauly (in press); Naamin (1984)

d/ Currie (1982)

e/ Thubtimsang and Meechukhan (1976); Thubtimsang (1977); Thubtimsaeng (1978, 1978a)

f/ Tiews, Bravo and Ronquillo (1972)

Table 5 (continued)

(2) Metapenaeus spp. See Sections 5 and 6.3 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY PENAEIDAE Metapenaeus affinis	Gulf of Thailand[a/]	10–19	–	–	–	2, 7	13 502–87 630	–	–	9.3–11.5	1:1	–	–	–	Molluscs, fish, crustaceans	M:W = $0.0000529L^{2.6182}$ F:W = $0.00000724L^{3.05}$
	South China Sea[b/] (Brunei)	10–45	3–4 g	19 g	Off shore to 45 m	2–4, 9–12	–	–	6–7	–	–	–	–	–	–	M:Log W = 2.939 Log C–0.036 F:Log W = 3.067 Log C–0.092
Metapenaeus brevicornis	Gulf of Thailand[c/]	–	5.0–9.5	–	–	All around	258 000–1 000 000	–	–	–	–	–	–	–	Phyto/zoo-planktons, feraminifera	
	South China Sea[b/] (Brunei)	–	–	–	2–4, 9–10	All around	–	1.7–5.2	All around	–	–	1 g/month	–	–	–	M:Log W = 3.124 Log C–0.091 F:Log W = 2.673 Log C–0.028
Metapenaeus ensis	Gulf of Thailand[d/]	10–19	–	–	–	1–4, 9–10	56 107–146 091	–	–	9.8	–	–	–	–	Cephalopods, fish, molluscs	M:W = $0.0002295L^{2.31}$ F:W = $0.00001737L^{2.858}$
	Java Sea[e/]	–	–	–	–	3.12	–	–	–	9.5 2.0 CL	–	–	–	–	–	–
Metapenaeus intermedius	Gulf of Thailand[f/]	20–39	13.4	19.4	–	3, 6, 9–12	45 200–604 359	–	–	9.0	1:1	–	–	–	Crustaceans, fish, cephalopods, molluscs, copepods, polychaeta	M:W = $0.000017L^{2.8534}$ F:W = $0.000023L^{2.858}$

a/ Thubtimsang and Meechukhan (1976); Thubtimsang (1977); Thubtimsaeng (1978, 1978a); Chaitiamvong and Thaiprayoon (1977, 1978a)

b/ Currie (1982)

c/ Chaitiamvong and Thaiprayoon (1979)

d/ Thubtimsang and Meechukhan (1976); Thubtimsang (1977); Chaitiamvong and Thaiprayoon (1979)

e/ Martosubroto (1977)

f/ Thubtimsang and Meechukhan (1976); Chaitiamvong and Phunwichian (1980)

Table 5 (2) (continued)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY PENAEIDAE Metapenaeus monoceros	Manila Bay[a/]	–	12.5	–	–	–	–	–	–	–	–	–	–	–	Zoo/phytoplanktons, benthic organisms, shrimps, annelida	–
	San Miguel Bay[a/]	–	–	–	–	–	–	–	–	–	–	–	–	–	Same as above	–

a/ Tiews, Bravo and Ronquillo (1972)

Table 5 (continued)

(3) Acetes spp. See Sections 5 and 6.3 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SERGESTIDAE Acetes erythraeus	Gulf of Thailand[a/]	–	0.8–3.3	3.3	–	–	–	–	–	–	0.78.1	–	–	–	–	–
Acetes indicus	Gulf of Thailand[a/]	–	1.4–2.6	3.3	–	–	–	–	–	–	0.6:1	–	–	–	–	–
Acetes japonicus	Gulf of Thailand[a/]	–	0.7–2.1	2.2	–	–	–	–	–	–	0.51:1	–	–	–	–	–
Acetes vulgaris	Gulf of Thailand[a/]	–	1.0–2.7	3.3	–	–	–	–	–	–	1.13:1	–	–	–	–	–

a/ Chaitianuong (1980a)

Table 6

Important biological features and parameters; cephalopods in the region
(1) Squids. See Sections 5 and 6.4 for annotations and broader references

(Body size refers to mantle length; sexes are combined unless specified as M: male or F: female)

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY LOLIGINIDAE Loligo edulis	Lombok[a/] Strait	10–20	–	–	–	–	–	–	–	–	–	–	–	–	–	$W=0.0066L^{1.74}$
Loligo duvaucellii	Gulf of Thailand[b/]	30–40	3.0–22.5	–	–	All around 2–5	1 000 000 4 000 000	–	–	5.6	1:1.3	–	–	–	–	M:W = $0.0081L^{1.189}$ F:W $0.0014L^{1.975}$
Loligo chinensis	Gulf of Thailand[c/]	40–50	4.5–40.9	40.9	–	All around 5, 11	800–7 000	–	–	–	–	–	–	–	Fish, molluscs, crustaceans	M:W = $0.0014L^{1.975}$ F:W = $0.000165L^{2.184}$
Loligo uyii	Gulf of Thailand[d/]	10–20	3.0–8.5	–	–	–	–	–	–	–	–	–	–	–	–	–
Sepietheutis lessoniana	Gulf of Thailand[e/]	20–30	3.0–26.1	26.1	–	All around	480–1 020	–	–	13.7	1:1.6	1.8–2.0 cm/month	–	–	Crustaceans, fish, cephalopods	M:W = $0.007657L^{1.846}$ F:W = $0.002437L^{2.10}$

a/ Marzuki (1983); Marzuki et al. (1983)

b/ Rattanaanun (1977, 1980)

c/ Rattanaanun (1977)

d/ Cholatarn (1978)

e/ Rattanaanun (1980)

Table 6 (continued)

(2) Cuttlefishes. See Sections 5 and 6.4 for annotations and broader references

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
			Mean (cm)	Maximum (cm)	Area	Season (month)		Size (cm)	Season (month)							
FAMILY SEPIIDAE Sepia pharaonis	Gulf of Thailand[a/]	20–30	3.8–26.0	26.0	–	All around 1, 7	780–2 500	–	–	14.3	1:1.8	1.5–2.0 cm/month	–	–	Crustaceans, fish	M:W = $0.000604L^{2.629}$ F:W = $0.000706L^{2.609}$
Sepia aculeata	Gulf of Thailand[b/]	10–30	2.0–18.9	–	–	All around 3–4, 7–9	650–3 900	–	–	8.1	1:1.4	–	–	–	Fish, crustaceans	M:W = $6.23\times10^{-4}L^{2.607}$ F:W = $4.88\times10^{-4}L^{2.677}$
Sepia recurvirostra	Gulf of Thailand[c/]	20–30	3.2–12.3	–	–	All around 2–3, 6–10	310–1 370	–	–	6.7	1:1.3	–	–	–	Crustaceans, fish, cephalopods	M:W = $0.001863L^{2.369}$ F:W = $0.00123L^{2.468}$
Sepia inermis	Gulf of Thailand[d/]	10–20	2.0–10.5	–	–	–	–	–	–	–	1:2.6	–	–	–	–	–
Sepia brevimana	Gulf of Thailand[a/]	30–40	2.4–8.2	9.0	–	–	3 312–6 565	–	–	–	1:2.2	–	–	–	Crustaceans, fish	M:F = $0.000607L^{2.599}$ W:F = $0.000764L^{2.549}$
Sepia lycidas	Gulf of Thailand[d/]	10–20	5.8–21.7	–	–	–	–	–	–	–	1:2	–	–	–	–	–

a/ Chotiyaputta (1982)

b/ Supongpan and Kongmuag (1976, 1976a)

c/ Supongpan and Kongmuag (1976, 1976a); Chotiyaputta (1982)

d/ Cholatarn (1978)

Table 6 (continued)

(3) Octopuses. See Sections 5 and 6.4 for annotations and broader references[a/]

Species	Area (country) surveyed	Vertical distribution range (m)	Body size captured		Spawning		Fecundity	Recruitment		Size at first maturity (cm)	Sex ratio (M:F)	Growth (rate or coefficient)	Mortality (coefficient)	Life span (year)	Food organisms	Length-weight relationship
FAMILY OCTOPODIDAE Octopus membranaceous	Gulf of Thailand[b/]	10-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Octopus dollfusi	Gulf of Thailand[b/]	10-50	-	-	-	-	-	-	-	-	-	-	-	-	-	-

a/ Although three other species are believed to be widely distributed in the region (Chikuni, 1983; Roper, Sweeney and Nauen, 1984), much attention has not been paid by fishermen

b/ Cholatarn (1978)

LIST OF FIGURES

		Page
1.	Definition of the sea area employed in this study	64
2.	Bathymetric condition of the region	65
3.	Geographic distribution; mackerels	66
4.	Geographic distribution; round scads	67
5.	Geographic distribution; sardines	68
6.	Geographic distribution; anchovies	69
7.	Geographic distribution; carangids	70
8.	Geographic distribution; king mackerels	71
9.	Geographic distribution; coastal tunas	72
10.	Geographic distribution; pony fish	73
11.	Geographic distribution; red snappers	74
12.	Geographic distribution; goat fish	75
13.	Geographic distribution; threadfin breams	76
14.	Geographic distribution; bigeye snappers	77
15.	Geographic distribution; groupers	78
16.	Geographic distribution; croakers	79
17.	Geographic distribution; lizardfish	80
18.	Geographic distribution; shrimps	81
19.	Geographic distribution; squids	82
20.	Geographic distribution; cuttlefish	83
21.	Estimated seasonal pattern of migration and displacement of major species groups in the region	84

Figure 1 Definition of the sea area employed in this study. See Section 3 of the text for annotations

Figure 2 Bathymetric condition of the region. See Section 3 of the text for annotations

Figure 3 Geographic distribution; mackerels. See Section 6.1.1 of the text for annotations and references

Figure 4 Geographic distribution; round scads. See Section 6.1.2 of the text for annotations and references

Figure 5 Geographic distribution; sardines. See Section 6.1.3 of the text for annotations and references

Figure 6 Geographic distribution; anchovies. See Section 6.1.4 of the text for annotations and references

Figure 7 Geographic distribution; carangids. See Section 6.1.5 of the text for annotations and references

Figure 8 Geographic distribution; king mackerels. See Section 6.1.6 of the text for annotations and references

Figure 9 Geographic distribution; coastal tunas. See Section 6.1.7 of the text for annotations and references

Figure 10 Geographic distribution; pony fish. See Section 6.2.1 of the text for annotations and references

Figure 11 Geographic distribution; red snappers. See Section 6.2.2 of the text for annotations and references

Figure 12 Geographic distribution; goat fish. See Section 6.2.3 of the text for annotations and references

Figure 13 Geographic distribution; threadfin breams. See Section 6.2.4 of the text for annotations and references

Figure 14 Geographic distribution; bigeye snappers. See Section 6.2.5 of the text for annotations and references

Figure 15 Geographic distribution; groupers. See Section 6.2.6 of the text for annotations and references

Figure 16 Geographic distribution; croakers. See Section 6.2.7 of the text for annotations and references

Figure 17 Geographic distribution; lizardfish. See Section 6.2.8 of the text for annotations and references

Figure 18 Geographic distribution; shrimps. See Section 6.3 of the text for annotations and references

Figure 19 Geographic distribution; squids. See Section 6.4 of the text for annotations and references

Figure 20 Geographic distribution; cuttlefish. See Section 6.4 of the text for annotations and references

ABBREVIATIONS
cp: coastal small pelagic fish
ma: mackerels, Rastrelliger spp.
sm: short mackerel, Rastrelliger brachysoma
rs: round scads, Decapterus spp.
sa: sardines, Sardinella spp.
os: Indian oil-sardine, Sardinella longiceps
an: anchovies, five genus included, most common, Stolephorus spp.
km: king mackerels, Scomberomorus spp.
ct: coastal tunas,
kawakawa, Euthynnus affinis
frigate tuna, Auxis thazard
longtail tuna, Thunnus tonggol
df: demersal fish
sn: snappers, Lutjanus and Priacanthus spp.
tr: threadfin bream, Nemipterus spp.
gr: groupers, Serranidae
cr: croakers, Sciaenidae
lz: lizardfish, Saurida spp.
sh: shrimps
Nautical miles
0 100 200 300 400
0 200 400 600 800
Kilometers

Figure 21 Estimated seasonal pattern of migration and displacement of major species groups in the region

Tipo-lito SAGRAF - Napoli

M-43
ISBN 92-5-102495-2